



(11) **EP 1 705 247 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 158(3) EPC

(43) Date of publication:
27.09.2006 Bulletin 2006/39

(21) Application number: **04799902.4**

(22) Date of filing: **25.11.2004**

(51) Int Cl.:
C12N 15/09 (2006.01) **C12N 1/15 (2006.01)**
C12N 1/19 (2006.01) **C12N 1/21 (2006.01)**
C12N 5/10 (2006.01) **C12N 9/02 (2006.01)**
C12P 17/08 (2006.01) **C07K 14/00 (2006.01)**

(86) International application number:
PCT/JP2004/017906

(87) International publication number:
WO 2005/052152 (09.06.2005 Gazette 2005/23)

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LU MC NL PL PT RO SE SI SK TR
Designated Extension States:
AL HR LT LV MK YU

(30) Priority: **27.11.2003 JP 2003396828**


(71) Applicants:
• **MERCIAN CORPORATION**
Chuo-ku,
Tokyo 104-8305 (JP)
• **Eisai Co., Ltd.**
Tokyo 112-0002 (JP)

(72) Inventors:
• **MACHIDA, Kazuhiro**
Iwata-shi,
Shizuoka (JP)
• **NAKASHIMA, Takashi**
Naka-gun, Kanagawa (JP)
• **ARITOKU, Yasuhide**
Iwata-shi, Shizuoka (JP)
• **TSUCHIDA, Toshio**
Iwata-shi, Shizuoka (JP)


(74) Representative: **HOFFMANN EITLE**
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

(54) **DNA PARTICIPATING IN HYDROXYLATION OF MACROLIDE COMPOUND**

(57) The present invention provides a DNA participating in biological transformation of a macrolide compound 11107B. The present invention provides, particularly, a DNA participating in biological transformation of a macrolide compound 11107B represented by the formula (I) into a 16-position hydroxy macrolide compound 11107D represented by the formula (II), the DNA encoding a protein having 16-position hydroxylating enzymatic activity or ferredoxin, to a method of isolating the DNA, to a protein encoded by the DNA, a plasmid carrying the DNA, a transformant obtained by transforming using the plasmid and a method of producing a 16-position hydroxy compound by using the transformant.



EP 1 705 247 A1




Description

Field of the invention


[0001] The present invention relates to a DNA participating in hydroxylation of a macrolide compound, a method of isolating it, a protein encoded by the DNA, a plasmid carrying the DNA, a transformant obtained by the transformation of the plasmid and a method of producing a 16-position hydroxy macrolide compound by using the transformant.

Prior Art

[0002] The 12-membered ring macrolide compound 11107D represented by the formula (II):



is a 12-membered ring macrolide compound having an excellent antitumor activity and has been found, together with a 12-membered ring macrolide compound 11107B represented by the formula (I):




from a cultured product of a *Streptomyces* sp. Mer-11107 strain (WO02/060890). The macrolide compound 11107D corresponds to a 16-position hydroxylated body of the macrolide compound 11107B. The productivity of the macrolide compound 11107D is lower than that of the macrolide compound 11107B and it has been therefore desired to establish an efficient method of producing the macrolide compound 11107D.

Disclosure of the invention


[0003] The purpose of the present invention is to find a DNA participating in hydroxylation of the macrolide compound 11107B to thereby provide a novel method of producing the macrolide compound 11107D.

[0004] The present invention relates to the following (1) to (15):

(1) a DNA participating in biological transformation of a macrolide compound (hereinafter referred to as a macrolide compound 11107B) represented by the formula (I):



15 into a 16-position hydroxy macrolide compound (hereinafter referred to as a macrolide compound 11107D) represented by the formula (II):



the DNA being an isolated and pure DNA comprising a DNA encoding a protein having 16-position hydroxylating enzymatic activity or ferredoxin, partly or entirely or its variant;

(2) the DNA described in (1), which is characterized by the following (a), (b) or (c):

(a) a DNA encoding a protein having the enzymatic activity in hydroxylating the 16-position of the macrolide compound 11107B and selected from the group consisting of a continuous nucleotide sequence from the base 1322 to base 2548 of the sequence No. 1; a continuous nucleotide sequence from the base 420 to base 1604 of the sequence No. 2; and a continuous nucleotide sequence from the base 172 to base 1383 of the sequence No. 3;

(b) a DNA which is a variant of the DNA described in the above (a);

(i) is hybridized with the DNA described in the above (a) under a stringent condition; and

(ii) encodes a protein having enzymatic activity in hydroxylating the 16-position of the macrolide compound 11107B; and

(c) a DNA encoding a protein having the same amino acid sequence as the protein encoded by the DNA described in the above (a) or (b) though it is not hybridized with the DNA described in the above (a) under a stringent condition because of the degeneracy of a gene codon;

(3) a protein encoded by the DNA as described in (2);

(4) a self-replicative or integrating replicative recombinant plasmid carrying the DNA as described in (2);

(5) a transformant into which the recombinant plasmid described in (4) transforms;

(6) a method of isolating a DNA encoding a protein having enzymatic activity in hydroxylating the 16-position of the macrolide compound 11107B, the method characterized by using the DNA as described in (2) or a DNA constituted of a part of the DNA as a probe or a primer;

(7) the DNA described in (1), which is characterized by the following (d), (e) or (f):

(d) a DNA encoding ferredoxin and selected from the group consisting of a continuous nucleotide sequence from the base 2564 to base 2761 of the sequence No. 1, a continuous nucleotide sequence from the base 1643 to base 1834 of the sequence No. 2 and a continuous nucleotide sequence from the base 1399 to base 1593 of the sequence No. 3;

(e) a DNA which is a variant of the DNA represented by the above (d);

(i) is hybridized with the DNA described in the above (d) under a stringent condition; and

(ii) encodes a protein having a ferredoxin function; and

(f) a DNA encoding a protein having the same amino acid sequence as the protein encoded by the DNA represented by the above (d) or (e) though it is not hybridized with the DNA described in the above (d) under a stringent condition because of the degeneracy of a gene codon;


(8) a protein encoded by the DNA as described in (7);

(9) a self-replicative or integrating replicative recombinant plasmid carrying the DNA as described in (7);


(10) a transformant into which the recombinant plasmid as described in (9) transforms;

(11) a method of isolating a DNA encoding a protein having a ferredoxin function, the method characterized by using the DNA as described in (7) or a DNA constituted of a part of the DNA as a probe or a primer;

(12) a method of producing a 16-position hydroxy macrolide compound, the method comprises the steps of culturing the transformant as described in (5) or (10) in a medium; bringing the proliferated transformant into contact with a macrolide compound represented by the formula (III):



(wherein W represents



[0005] R¹², R^{16b}, R^{17a}, R^{17b}, R¹⁸, R^{20a}, R^{20b}, R^{21a} and R^{21b}, which may be the same as or different from, respectively represent:

(1) hydrogen atom;

(2) a C₁₋₂₂ alkyl group which may have a substituent;

(3) -OR (wherein R represents:

1) hydrogen atom; or

2) a C₁₋₂₂ alkyl group;

3) a C₇₋₂₂ aralkyl group;

4) a 5-membered to 14-membered heteroaryloxyalkyl group;

5) a C₂₋₂₂ alkanoyl group;

6) a C₇₋₁₅ aroyl group;

7) a C₃₋₂₃ unsaturated alkanoyl group;

8) -COR^{co} (wherein R^{co} represents:

8-1) a 5-membered to 14-membered heteroaryloxyaryl group;

- 8-2) a C₁₋₂₂ alkoxy group;
 8-3) an unsaturated C₂₋₂₂ alkoxy group;
 8-4) a C₆₋₁₄ aryloxy group;
 8-5) a 5-membered to 14-membered heteroaryloxy group; or
 8-6) a 3-membered to 14-membered nitrogen-containing non-aromatic heterocyclic group, each of which may have a substituent);

- 9) a C₁₋₂₂ alkylsulfonyl group;
 10) a C₆₋₁₄ arylsulfonyl group; or
 11) -SiR^{s1}R^{s2}R^{s3}, (wherein R^{s1}, R^{s2} and R^{s3}, which may be the same as or different from, respectively represent a C₁₋₆ alkyl group or a C₆₋₁₄ aryl group), each of which may have a substituent);

(4) a halogen atom; or

(5) -R^M-NR^{N1}R^{N2}, {wherein R^M represents a single bond or -O-CO-; and R^{N1} and R^{N2}

1) may be the same as or different from, respectively represent:

- 1-1) hydrogen atom; or
 1-2)

- (i) a C₁₋₂₂ alkyl group;
 (ii) an unsaturated C₂₋₂₂ alkyl group;
 (iii) a C₂₋₂₂ alkanoyl group;
 (iv) a C₇₋₁₅ aroyl group;
 (v) an unsaturated C₃₋₂₃ alkanoyl group;
 (vi) a C₆₋₁₄ aryl group;
 (vii) a 5-membered to 14-membered heteroaryl group;
 (viii) a C₇₋₂₂ aralkyl group;
 (ix) a C₁₋₂₂ alkylsulfonyl group; or
 (x) a C₆₋₁₄ arylsulfonyl group, each of which may have a substituent, or


2) and R^{N1} and R^{N2} may be combined with the nitrogen atom to which they bound, to form a 3-membered to 14-membered nitrogen-containing non-aromatic heterocyclic group}, provided that

R^{21a} and R^{21b} may be combined with each other to form (i) a ketone structure (=O) or (ii) an oxime structure {=NOR^{ox} (wherein R^{ox} represents a C₁₋₂₂ alkyl group, an unsaturated C₂₋₂₂ alkyl group, a C₆₋₁₄ aryl group, a 5-membered to 14-membered heteroaryl group or a C₇₋₂₂ aralkyl group, each of which may have a substituent))};

R^{16a} represents hydrogen atom;


R^{21c} represents:

- (1) hydrogen atom; or
 (2)




(wherein R^{22a}, R^{22b} and R^{22c}, which may be the same as or different from, respectively represent:

- 1) hydrogen atom;
 2) a C₁₋₆ alkyl group;
 3) -OR (wherein R has the same meaning as the above);
 4) -R^M-NR^{N1}R^{N2} (wherein R^M, R^{N1} and R^{N2} have the same meanings as the above); or
 5) a halogen atom, or any one of R^{21a} and R^{21b} may be combined with any one of R^{22a} and R^{22b} to form the partial structure;



); and G^m represents:

(1) a group represented by the formula (GM-I):



{wherein

R^2 and R^{10} , which may be the same as or different from, respectively represent hydrogen atom or a C_{1-22} alkyl group;

R^{3a} , R^{3b} , R^{5a} , R^{5b} , R^{6a} and R^{6b} , which may be the same as or different from, respectively represent:

- 1) hydrogen atom;
- 2) hydroxyl group;
- 3)

3-1) a C_{1-22} alkyl group;

3-2) a C_{1-22} alkoxy group;

3-3) a C_{6-14} aryloxy group;

3-4) a 5-membered to 14-membered heteroaryloxy group;

3-5) a C_{2-22} alkanoyloxy group;

3-6) a C_{7-15} aroyloxy group;

3-7) a C_{3-23} unsaturated alkanoyloxy group;

3-8) $-OCOR^{co}$ (wherein R^{co} has the same meaning as the above);

3-9) a C_{1-22} alkylsulfonyloxy group;

3-10) a C_{6-14} arylsulfonyloxy group; or

3-11) $-OSiR^{s1}R^{s2}R^{s3}$ (wherein R^{s1} , R^{s2} and R^{s3} have the same meanings as the above), each of which may have a substituent;

4) a halogen atom; or


5) $-R^M-NR^{N1}R^{N2}$ (wherein R^M , R^{N1} and R^{N2} have the same meanings as the above); or

R^{5a} and R^{5b} may be combined with each other to form a ketone structure ($=O$); or


R^{6a} and R^{6b} may be combined with each other to form a spirooxysilanyl group or an exomethylene group; or

R^{7a} and R^{7b} , which may be the same as or different from, respectively represent hydrogen atom or $-OR^H$ (wherein R^H represents hydrogen atom, a C_{1-22} alkyl group or a C_{2-22} alkanoyl group));


(2) a group represented by the formula (GM-II):



15 (wherein R^2 , R^{3a} , R^{3b} , R^{6a} , R^{6b} , R^{7a} , R^{7b} and R^{10} have the same meanings as those in the formula (GM-I));
(3) a group represented by the formula (GM-III):




30 (wherein R^2 , R^{5a} , R^{5b} , R^{6a} , R^{6b} , R^{7a} , R^{7b} and R^{10} have the same meanings as those in the formula (GM-I));
(4) a group represented by the formula (GM-IV):




45 (wherein R^2 , R^{6a} , R^{7a} , R^{7b} and R^{10} have the same meanings as those in the formula (GM-I)); or
(5) a group represented by the formula (GM-V):

50

55




wherein R^2 , R^{3a} , R^{6a} , R^{6b} and R^{10} have the same meanings as those in the formula (GM-I)) during or after culturing, to convert it into a 16-position hydroxy macrolide compound represented by the formula (IV):



(wherein W , R^{12} , R^{16b} , R^{17a} , R^{17b} , R^{20a} , R^{20b} , R^{21a} , R^{21b} , R^{21c} and G^m have the same meanings as those in the formula (III)); and then collecting the 16-position hydroxy macrolide compound thus converted;

(13) a production method according to (12), wherein the transformant is the transformant as described in (5) and has a DNA encoding ferredoxin;

(14) the production method as described in (12), the method comprises the step of converting a compound represented by the formula (III-a) :




(wherein



represents a double bond or a single bond; W' represents a double bond or



; R^{5'} represents hydrogen atom or an acetoxy group; R^{6'} represents hydrogen atom or hydroxyl group; and R^{7'} represents hydrogen atom or acetyl group) into a compound represented by the formula (IV-a):



(wherein




, W', R^{5'}, R^{6'} and R^{7'} have the same meanings as those in the formula (III-a));

(15) the production method as described in (14), wherein, in the conversion of the compound of the formula (III-a) into the compound of the formula (IV-a), the compound to be subjected is a compound selected from the group consisting of:

(1) a compound in which



is a single bond; W' is



and R^{5'}, R^{6'} and R^{7'} are respectively hydrogen atom;

(2) a compound in which



is a single bond, W' is



R^{5'} and R^{6'} are respectively hydrogen atom; and R^{7'} is acetyl group;

(3) a compound in which



is a single bond, W' is




R^{5'} and R^{7'} are respectively hydrogen atom; and R^{6'} is hydroxyl group;

(4) a compound in which



is a single bond, W' is



R^{5'} is hydrogen atom, R^{6'} is hydroxy group; and R^{7'} is acetyl group;

(5) a compound in which



is a single bond; W' is a double bond; and R^{5'}, R^{6'} and R^{7'} are respectively hydrogen atom;

(6) a compound in which



is a single bond; W' is a double bond; R^{5'} and R^{6'} are respectively hydrogen atom; and R^{7'} is acetyl group;

(7) a compound in which



is a single bond; W' is a double bond; R^{5'} and R^{7'} are respectively hydrogen atom; and R^{6'} is hydroxyl group;

(8) a compound in which



is a single bond; W' is a double bond; R^{5'} is hydrogen atom; R^{6'} is hydroxy group; and R^{7'} is acetyl group;

(9) a compound in which



is a double bond; W' is



R^{5'} and R^{7'} are respectively hydrogen atom; and R^{6'} is hydroxyl group;

(10) a compound in which



is a double bond; W' is



R^{5'} is hydrogen atom; R^{6'} is hydroxy group; and R^{7'} is acetyl group;

(11) a compound in which



is a single bond; W' is




R^{5'} is acetoxy group; R^{6'} is hydroxyl group; and R^{7'} is hydrogen atom; and

(12) a compound in which



is a single bond; W' is



R^{5'} is an acetoxy group; R^{6'} is hydroxyl group; and R^{7'} is acetyl group; and

(16) use of the transformant as described in (5) or (10) for producing a 16-position hydroxy macrolide compound.

[0006] The present invention made it possible to isolate a DNA encoding a protein having the enzymatic activity in hydroxylating the 16-position of a macrolide compound 11107B or ferredoxin and to determine its nucleotide sequence. Moreover, a plasmid carrying the DNA and a transformant into which the plasmid transformed were formed and a 16-position hydroxy macrolide compound could be produced using the transformant in an efficient manner.

[0007] Hereinafter, embodiments of the present invention will be explained in detail.

Microorganisms having the ability of converting a macrolide compound 11107B into a macrolide compound 11107D

[0008] In the present invention, a DNA encoding a protein having enzymatic activity in hydroxylating the 16-position or ferredoxin, partly or entirely can be isolated from the mycelia isolated from a culture broth in which microorganisms having the ability of converting the macrolide compound 11107B into the macrolide compound 11107D are cultured and the nucleotide sequence of the DNA can be determined. Then, a self-replicative or integrating replicative recombinant plasmid carrying the DNA is architecturally formed and a transformant is prepared using the plasmid.

[0009] The transformant thus prepared is cultured in the culture media and the proliferated transformant is brought into contact with the macrolide compound represented by the above formula (III) during or after culturing, to thereby convert the macrolide compound into the 16-position hydroxy macrolide compound represented by the formula (IV) and the converted 16-position hydroxy macrolide compound is collected, whereby the 16-position hydroxy macrolide compound can be obtained.

[0010] Any microorganisms having the ability of converting the macrolide compound 11107B into the macrolide compound 11107D may be used irrespective of the type of species and strain. Preferable examples of the microorganisms may include a Streptomyces sp. Mer-11107 or A-1544 strain and an unidentified Actinomyces A-1560 strain which were each isolated from soils.

[0011] It is to be noted that the Streptomyces sp. Mer-11107 was deposited as FERM P-18144 at the National Institute of Bioscience and Human-Technology Agency of Industrial Science and Technology (1-3, Higashi 1-chome Tsukuba-

shi, Ibaraki-ken 305-8566 Japan) as of December 19, 2000, and then transferred to International Deposit FERM BP-7812 at International Patent Organism Depository (IPOD) National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1, Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken 305-8566 Japan) as of November 27, 2001. The A-1544 strain was deposited as **FERM P-18943** at International Patent Organism Depository National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1, Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken 305-8566 Japan) as of July 23, 2002, and then transferred to International Deposit FERM BP-8446 as of July 30, 2003, at International Patent Organism Depository (IPOD) National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1, Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken 305-8566 Japan). The A-1560 strain was deposited as FERM P-19585 at the National Institute of Bioscience and Human-Technology Agency of Industrial Science and Technology (1-3, Higashi 1-chome Tsukuba-shi, Ibaraki-ken 305-8566 Japan) as of November 13, 2003 and then transferred to International Deposit FERM BP-10102 as of August 19, 2004, at International Patent Organism Depository (IPOD) National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1, Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken 305-8566 Japan).

[0012] The taxonomical properties of the above strains are as follows.

(Taxonomical properties of the Mer-11107 strain)

(1) Morphological characteristics

[0013] Spirilles type aerial hyphae were extended from the vegetative hyphae. Spore chains consisting of about 10 to 20 cylindrical spores were formed at the end of the matured aerial hyphae. The size of the spores was about $0.7 \times 1.0 \mu\text{m}$, the surface of the spores was smooth, and specific organs such as sporangium, sclerotium and flagellum were not observed.

(2) Cultural characteristics on various media

[0014] Cultural characteristics of the strain after incubation at 28°C for two weeks on various media are shown as follows. The color tone is described by the color names and codes which are shown in the parentheses of Tresner's Color wheels.

1) Yeast extract-malt extract agar medium

The strain grew well, the aerial hyphae grew up on the surface, and light gray spores (Light gray; d) were observed. The reverse side of colony was Light melon yellow (3ea). Soluble pigment was not produced.

2) Oatmeal agar medium

The strain grew moderately, the aerial hyphae grew slightly on the surface, and gray spores (Gray; g) were observed. The reverse side of colony was Nude tan (4gc) or Putty (1 1/2ec). Soluble pigment was not produced.

3) Inorganic salts-starch agar medium

The strain grew well, the aerial hyphae grew up on the surface, and gray spores (Gray; e) were observed. The reverse side of colony was Fawn (4ig) or Gray (g). Soluble pigment was not produced.

4) Glycerol-asparagine agar medium

The strain grew well, the aerial hyphae grew up on the surface, and white spores (White; a) were observed. The reverse side of colony was Pearl pink (3ca). Soluble pigment was not produced.

5) Peptone-yeast extract-iron agar medium

The strain growth was bad, and the aerial hyphae did not grow on the surface. The reverse side of colony was Light melon yellow (3ea). Soluble pigment was not produced.

6) Tyrosine agar medium

The strain grew well, the aerial hyphae grew up on the surface, and white spores (White; a) were observed. The reverse side of colony was Pearl pink (3ca). Soluble pigment was not produced.

(3) Utilization of various carbon sources

[0015] Various carbon sources were added to Pridham-Gottlieb agar and incubated 28°C for 2 weeks. The growth of the strain is shown below.

1) L-arabinose ±

2) D-xylose ±

3) D-glucose +

4) D-fructose +

- 5) Sucrose +
 6) Inositol +
 7) L-rhamnose -
 8) D-mannitol +
 9) Raffinose +
 (+ positive, ± slightly positive, - negative)

(4) Various physiological properties

[0016] Various physiological properties of the present strain are as follows.

- (a) Range of growth temperature (yeast extract-malt extract agar, incubation for 2 weeks): 12°C to 37°C
 (b) Range of optimum growth temperature (yeast extract-malt extract agar, incubation for 2 weeks): 21°C to 33°C
 (c) Liquefaction of gelatin (glucose-peptone-gelatin medium): negative
 (d) Coagulation of milk (skim milk medium): negative
 (e) Peptonization of milk (skim milk medium): negative
 (f) Hydrolysis of starch (inorganic salts-starch agar) : positive (g) Formation of melanoid pigment (peptone-yeast extract-iron agar): negative
 (tyrosine agar): negative
 (h) Production of hydrogen sulfide (peptone-yeast extract-iron agar): negative
 (i) Reduction of nitrate (broth containing 0.1% potassium nitrate): negative
 (j) Sodium chloride tolerance (yeast extract-malt extract agar, incubation for 2 weeks): grown at a salt content of 4% or less

(5) Chemotaxonomy

[0017] LL-diaminopimelic acid and glycine were detected from the cell wall of the present strain.
 (Taxonomical properties of the A-1544 strain)

(1) Morphological characteristics

[0018] Spira type aerial hyphae were extended from vegetative hyphae in this strain. Spore chains consisting of about 10 to 20 of cylindrical spores were formed at the end of the matured aerial hyphae. The size of the spores was about 1.0 x 1.2 to 1.4 µm, the surface of the spores was spiny, and specific organs such as sporangium, sclerotium and flagellum were not observed.

(2) Cultural characteristics on various media

[0019] Cultural characteristics of the strain after incubation at 28°C for two weeks on various media are shown in Table 1. The color tone is described by the color names and codes which are shown in the parentheses of Tresner's Color wheels.

Table 1

Medium	Growth	Aerial hyphae	Color of vegetative hyphae	Soluble pigment
Yeast extract -malt extract agar (ISP-2)	Good	Thick Silver gray (3fe)	Light melon yellow (3ea)	None
Oatmeal agar (ISP-3)	Good	Abundant Light gray to Silver gray (d to 3fe)	Light melon yellow (3ea)	None
Inorganic salts - starch agar (ISP-4)	Good	Abundant Silver gray (3fe)	Light melon yellow (3ea)	None
Glycerol - asparagine agar (ISP-5)	Good	Abundant Ashes (5fe)	Light melon yellow (3ea)	None
Peptone-yeast extract - iron agar (ISP-6)	Good	None	Light melon yellow (3ea)	Pale blackish brown

(continued)

Medium	Growth	Aerial hyphae	Color of vegetative hyphae	Soluble pigment
Tyrosine agar (ISP-7)	Good	Abundant Covert gray (2fe)	Light melon yellow (3ea)	None

(3) Utilization of various carbon sources

[0020] Various carbon sources were added to Pridham-Gottlieb agar and incubated at 28°C for 2 weeks. The growth of the strain is shown in Table 2.

Table 2

D-glucose	+	inositol	-
L-arabinose	+	L-rhamnose	+
D-xylose	+	D-mannitol	+
D-fructose	+	raffinose	-
sucrose	-		
+: positive, ±: slightly positive, -: negative			

(4) Various physiological properties

[0021] Various physiological properties of the present strain are as follows.

- (a) Range of growth temperature (yeast extract-malt extract agar, incubation for 2 weeks): 15°C to 41°C
- (b) Range of optimum growth temperature (yeast extract-malt extract agar, incubation for 2 weeks): 20°C to 37°C
- (c) Liquefaction of gelatin (glucose-peptone-gelatin medium) : positive
- (d) Coagulation of milk (skim milk medium): positive
- (e) Peptonization of milk (skim milk medium): positive
- (f) Hydrolysis of starch (inorganic salts-starch agar): positive
- (g) Formation of melanoid pigment (peptone-yeast extract-iron agar): positive (tyrosine agar): negative
- (h) Production of hydrogen sulfide (peptone-yeast extract-iron agar): positive
- (i) Reduction of nitrate (broth containing 0.1% potassium nitrate): negative
- (j) Sodium chloride tolerance (yeast extract-malt extract agar, incubation for 2 weeks) : grown at a salt content of 7% or less

(5) Chemotaxonomy

[0022] LL-diaminopimelic acid was detected from the cell wall of the present strain.

DNA of the present invention

[0023] The present inventors have isolated a DNA participating in the hydroxylation of the 16-position of a macrolide compound, specifically, a DNA encoding a protein having 16-position hydroxylating enzymatic activity and a DNA encoding a protein having a ferredoxin function from the above microorganisms and determined the nucleotide sequence of the DNA. The DNA encoding a protein having 16-position hydroxylating enzymatic activity and the DNA encoding a protein having a ferredoxin function are hereinafter generically called "a 16-position hydroxylating enzyme relevant DNA" as the case may be.

[0024] The DNA encoding a protein having 16-position hydroxylating enzymatic activity is those represented by the following (1-1), (1-2) or (1-3):

- (1-1) a DNA selected from those having a continuous nucleotide sequence from the base 1322 to base 2548 of the sequence No. 1, a continuous nucleotide sequence from the base 420 to base 1604 of the sequence No. 2 and a

continuous nucleotide sequence from the base 172 to base 1383 of the sequence No. 3;
 (1-2) a DNA which is a variant of the DNA described in the above (1-1);

- (i) is hybridized with any one of the DNAs described in the above (1-1) under a stringent condition; and
 (ii) codes a protein having enzymatic activity in hydroxylating the 16-position of the macrolide compound; and

(1-3) a DNA encoding a protein having the same amino acid sequence as the protein encoded by the DNA described in the above (1-1) or (1-2) though it is not hybridized with any of the DNA described in the above (1-1) or (1-2) under a stringent condition because of the degeneracy of a gene codon.

[0025] The "16-position hydroxylating enzymatic activity" means such enzymatic activity as to hydroxylate the 16-position of the macrolide compound 11107B represented by the formula (I) to thereby convert the macrolide compound into the macrolide compound 11107D represented by the formula (II).

[0026] The DNA encoding a protein having a ferredoxin function in the present invention is those represented by the following (2-1), (2-2) or (2-3).

(2-1) a DNA encoding ferredoxin and selected from the group consisting of a continuous nucleotide sequence from the base 2564 to base 2761 of the sequence No. 1, a continuous nucleotide sequence from the base 1643 to base 1834 of the sequence No. 2 and a continuous nucleotide sequence from the base 1399 to base 1593 of the sequence No. 3;

(2-2) a DNA which is a variant of the DNA described in the above (2-1);

- (i) is hybridized with the DNA described in the above (2-1) under a stringent condition; and
 (ii) codes a protein having a ferredoxin function; and

(2-3) a DNA encoding a protein having the same amino acid sequence as the protein encoded by the DNA represented by the above (2-1) or (2-2) though it is not hybridized with the DNA represented by the above (2-1) under a stringent condition because of the degeneracy of a gene codon.

[0027] "The ferredoxin function" means the protein function of transferring electrons to the above 16-position hydroxylating enzyme to bear a role together with the above 16-position hydroxylating enzyme in the hydroxylation reaction.

[0028] Also, "the nucleotide sequence hybridized under a stringent condition" means a DNA nucleotide sequence obtained when any one of the DNAs of the above (1-1) and (1-2) is used as a probe and, for example, a colony hybridization method, plaque hybridization method or Southern blot hybridization method is used. Examples of the DNA having such a nucleotide sequence may include those identified by carrying out hybridization in the presence of 0.7 to 1.0 M NaCl at 65°C using a filter to which a DNA derived from a colony or a plaque or a fragment of the DNA is fixed and then washing the filter at 65°C by using 0.1 to 2 × SSC solution (1 × SSC solution: 150 mM sodium chloride and 15 mM sodium citrate). The hybridization may be carried out according to the method described in Molecular Cloning: A laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY., 1989 (hereinafter abbreviated as molecular cloning, 2nd ed.).

[0029] Examples of the DNA hybridized in a stringent condition include DNAs having a nucleotide sequence having a certain level or more of homology with the nucleotide sequence of the DNA to be used as the probe, and specifically DNAs having a nucleotide sequence having 80% or more, preferably 85% or more, more preferably 90% or more, still more preferably 95% or more and most preferably 98% or more homology with the nucleotide sequence of the DNA used as the probe.

[0030] There is no particular limitation to a method of obtaining the 16-position hydroxylating enzyme relevant DNA. An appropriate probe or a primer is prepared based on the information of the nucleotide sequence described in the sequence No. 1, No. 2 or No. 3 of the sequence chart in this specification. Using the probe or primer, a DNA library of microorganisms belonging to Actinomyces is screened, and thus the DNA of the present invention can be isolated. The DNA library can be produced by the usual method from microorganisms expressing the aforementioned 16-position hydroxylating enzymatic activity.

[0031] The 16-position hydroxylating enzyme relevant DNA of the present invention can also be obtained by a PCR method. A DNA library derived from the aforementioned microorganisms is used as a template and a pair of primers which are so designed as to amplify any one of the nucleotide sequences described in the sequence No. 1, No. 2 or No. 3 are used to carry out PCR. The reaction condition of the PCR may be appropriately designed. Examples of the reaction condition may include the condition of a process in which the cycle of the process involving a reaction run at 94°C for 30 seconds (denaturing), a reaction run at 55°C for 30 seconds to one minute (annealing) and a reaction run at 72°C for 2 minutes (extension) is repeated 30 times and then a reaction is run at 72°C for 7 minutes. Then, the

amplified DNA fragment can be cloned in a vector which can be amplified in a proper host.

[0032] The aforementioned operations such as the preparation of a probe or a primer, the construction of a DNA library, the screening of a DNA library and the cloning of a target gene are obvious to a person skilled in the art and may be carried out according to methods as described in for example, Molecular Cloning 2nd ed., Current Protocols in Molecular Biology, Supplement 1 to 38, John Wiley & Sons (1987-1997).

[0033] No particular limitation is imposed on a method of obtaining the protein in the present invention. The protein may be a protein synthesized by chemical synthesis or a recombinant protein produced by gene recombination techniques. When the recombinant protein is produced, first, the DNA encoding the protein as described above in this specification is obtained. The protein of the present invention can be produced by introducing this DNA into a proper expression system. Manifestation of the protein in the expression system will be described later in the specification.

Recombinant vector in the present invention

[0034] The DNA of the present invention may be used in the situation where it is inserted in an appropriate vector. No particular limitation is imposed on the kind of the vector to be used in the present invention and the vector may be either a self-replicative one (for example, a plasmid) or one that is incorporated into a genome of a host cell when introduced into the host cell and is replicated together with the incorporated chromosome. In the expression vector, the DNA of the present invention is operationally linked to elements (for example, a promoter) which are necessary for transcription. The promoter is a DNA sequence exhibiting transcriptional activity in a host cell and may be selected suitably corresponding to the type of host.

The transformant of the present invention and production of a recombinant protein using the transformant

[0035] The transformant may be produced by introducing the DNA or recombinant vector of the present invention into an appropriate host. The host cell into which the DNA or recombinant vector of the present invention is introduced may be any desired cell which can express the gene according to the present invention. Examples of the host cell include bacteria, yeast, fungi and higher eucaryote cells. Examples of the bacterial cell include Gram-positive bacteria such as *Bacillus* or *Streptomyces* or Gram-negative bacteria such as *E. coli*. The transformation of these bacteria may be accomplished using a competent cell according to a protoplast method, electroporation method or other known methods. For example, the electroporation method may be performed as follows. A plasmid into which a foreign gene is inserted is added to a suspension of the competent cell, this suspension is poured into a cuvet specially used for an electroporation method and high-voltage electric pulse is applied to the cuvet. Then, the cells are cultured in a selective medium and a transformant is isolated on a plate agar media.

[0036] Examples of the yeast cell include cells belonging to *Saccharomyces* or *Schizosaccharomyces*. Specific examples of the yeast cell include *Saccharomyces cerevisiae* or *Saccharomyces kluyveri*. Examples of a method of introducing the recombinant vector into the yeast host may include an electroporation method, spheroplast method and lithium acetate method. Examples of the above other fungus cell include mycotic cells belonging to *Aspergillus*, *Neurospora*, *Fusarium* or *Trichoderma*. When mold fungi are used as the host cell, a DNA architecture is incorporated into a host chromosome to obtain a recombinant host cell, whereby transformation can be accomplished. The incorporation of the DNA architecture into the host chromosome can be accomplished, for example, by homologous recombination or heterologous recombination.

[0037] The above transformant is cultured in an appropriate nutrient medium under the condition enabling the expression of the introduced gene. In order to isolate the protein of the invention from the culture product of the transformant and to purify the protein, the usual protein isolating and purifying method may be used.

[0038] For example, when the protein of the present invention is expressed in a soluble form in cells, the cells are collected by centrifugation after the cultivation is finished and are suspended in a buffer solution. Then, the suspended solution is subjected to, for example, a ultrasonic crusher to break the cells, thereby obtaining a cell-free extract and the cell-free extract is centrifuged. A purified preparation can be obtained from the obtained supernatant by combining measures such as the usual isolation and purifying methods such as a solvent extraction method, salting-out method using ammonium sulfate, desalting method, precipitation method using an organic solvent, anion exchange chromatography using a resin such as diethylaminoethyl (DEAE) sepharose, cation exchange chromatography using a resin such as SP-Sephacrose FF (manufactured by Amasham Bioscience Company), hydrophobic chromatography using a resin such as butyl sepharose and phenyl sepharose, gel filtration method using a molecular sieve, affinity chromatography, chromatofocusing method and electrophoresis method such as an isoelectric focusing electrophoresis.

Method of production of 16-position hydroxy macrolide compounds

[0039] The present invention involves a method of producing 16-position hydroxy macrolide compounds represented by the above formula (IV), the method comprising using a transformant into which a DNA encoding a protein having 16-position hydroxylating enzymatic activity or a protein having a ferredoxin function is introduced and hydroxylating mac-

rolide compounds represented by the above formula (III) in the presence of the transformant.

[0040] The macrolide compounds that can be hydroxylated by the transformant of the present invention is macrolide compounds represented by the above formula (III) (macrolide compounds represented by the above formula (IV)), preferably macrolide compounds represented by the above formula (III-a) (macrolide compounds represented by the above formula (IV-a) and more preferably the macrolide compound 11107B (macrolide compound 11107D). The compounds in the parenthesis are 16-position hydroxy macrolide compounds that are hydroxylated products.

[0041] The condition under which the macrolide compounds are hydroxylated in the presence of the transformant is as follows.

[0042] First, the 16-position hydroxylating enzyme relevant DNA in the transformant is expressed by adding, if necessary, inducing materials. The strain expressing the DNA is brought into contact with the macrolide compounds represented by the above formula (III) to run a conversion reaction. The temperature of the conversion reaction may be suitably determined taking the optimum growth temperature of the transformant into account. The reaction time may also be suitably determined in consideration of the conversion rate (degree of progress of the reaction) into the 16-position hydroxy macrolide compound. For example, the condition of 20 to 31°C and 1 to 5 days is preferable. Moreover, as to the reaction system, the reaction may be run in any system including a batch system or a continuous system.

[0043] For the isolation and purifying of the produced 16-position hydroxymacrolide compounds, the separation and purifying method used usually to isolate a microbial metabolite from the culture broth may be utilized. All known separation and purifying methods such as organic solvent extraction using methanol, ethanol, acetone, butanol, ethyl acetate, butyl acetate, chloroform or toluene, absorption chromatograph using a hydrophobic adhesive resin such as Diaion HP-20, gel filtration chromatography using Sefadex LH-20, adsorption chromatography using activated carbon, silica gel or the like, absorption chromatograph using thin-layer chromatography and high-performance liquid chromatography using an reverse phase column are equivalent to these separation and purifying methods. The separation and purifying method is not limited to these methods shown here. These methods may be used singly or in combinations of two or more in an optional order or repeatedly, which makes it possible to isolate and purify the target 16-position hydroxy macrolide compounds.

[0044] The variant of the DNA in the present invention means a DNA that is obtained by modifying the DNA by deletion, conversion, addition or insertion treatments in the structural base of the DNA or its derivatives and shows the same effects as the original DNA.

[0045] The "halogen atom" used in the specification of the present application means a fluorine atom, a chlorine atom, a bromine atom and an iodine atom.

[0046] The " C_{1-22} alkyl group" used in the specification of the present application indicates a linear or branched alkyl group having 1 to 22 carbon atoms, such as methyl group, ethyl group, n-propyl group, iso-propyl group, n-butyl group, iso-butyl group, sec-butyl group, tert-butyl group, n-pentyl group, 1,1-dimethylpropyl group, 1,2-dimethylpropyl group, 2,2-dimethylpropyl group, 1-ethylpropyl group, n-hexyl group, 1-ethyl-2-methylpropyl group, 1,1,2-trimethylpropyl group, 1-methylbutyl group, 2-methylbutyl group, 1,1-dimethylbutyl group, 1,2-dimethylbutyl group, 2,2-dimethylbutyl group, 1,3-dimethylbutyl group, 2,3-dimethylbutyl group, 1-ethylbutyl group, 2-ethylbutyl group, 2-methylpentyl group, 3-methylpentyl group, n-heptyl group, n-octyl group, n-nonyl group or n-decyl group; preferably a linear or branched alkyl group having 1 to 6 carbon atoms, such as methyl group, ethyl group, n-propyl group, iso-propyl group, n-butyl group, iso-butyl group, sec-butyl group or tert-butyl group.

[0047] The "unsaturated C_{2-22} alkyl group" used in the specification of the present application indicates a linear or branched alkenyl group having 2 to 22 carbon atoms or a linear or branched alkynyl group having 2 to 22 carbon atoms, such as vinyl group, allyl group, 1-propenyl group, isopropenyl group, 2-methyl-1-propenyl group, 2-methyl-2-propenyl group, 1-butenyl group, 2-butenyl group, 3-butenyl group, 1-pentenyl group, 1-hexenyl group, 1,3-hexanedieryl group, 1,5-hexanedieryl group, ethynyl group, 1-propynyl group, 2-propynyl group, 1-butylnyl group, 2-butylnyl group, 3-butylnyl group, 1-ethynyl-2-propynyl group, 2-methyl-2-propynyl group, 1-pentynyl group, 1-hexynyl group, 1,3-hexanediynyl group or 1,5-hexanediynyl group. It preferably indicates a linear or branched alkenyl group having 2 to 10 carbon atoms or a linear or branched alkynyl group having 2 to 10 carbon atoms, such as vinyl group, allyl group, 1-propenyl group, isopropenyl group, ethynyl group, 1-propynyl group, 2-propynyl group, 1-butylnyl group, 2-butylnyl group or 3-butylnyl group.

[0048] The " C_{6-14} aryl group" used in the specification of the present application means an aromatic hydrocarbon group having 6 to 14 carbon atoms, and a monocyclic group and condensed rings such as a bicyclic group and a tricyclic group are included. Examples thereof are phenyl group, indenyl group, 1-naphthyl group, 2-naphthyl group, azulenyl group, heptalenyl group, indacenyl group, acenaphthyl group, fluorenyl group, phenalenyl group, phenanthrenyl group and anthracenyl group; of which a preferred example is phenyl group, 1-naphthyl group or 2-naphthyl group.

[0049] The "5-membered to 14-membered heteroaryl group" used in the specification of the present application means a monocyclic, bicyclic or tricyclic 5-membered to 14-membered aromatic heterocyclic group which contains one or more of hetero atoms selected from the group consisting of a nitrogen atom, sulfur atom and oxygen atom. Preferred examples thereof are a nitrogen-containing aromatic heterocyclic group such as pyrrolyl group, pyridinyl group, pyridazinyl group,

pyrimidinyl group, pyrazinyl group, triazolyl group, tetrazolyl group, benzotriazolyl group, pyrazolyl group, imidazolyl group, benzimidazolyl group, indolyl group, isoindolyl group, indoliziny group, purinyl group, indazolyl group, quinolinyl group, isoquinolinyl group, quinoliziny group, phthalazinyl group, naphthyridinyl group, quinoxalinyl group, quinoxalinyl group, cinnolinyl group, pteridinyl group, imidazotriazinyl group, pyrazinopyridazinyl group, acridinyl group, phenanthridinyl group, carbazolyl group, carbazolinyl group, perimidinyl group, phenanthrolinyl group, phenazinyl group, imidazopyridinyl group, imidazopyrimidinyl group, pyrazolopyridinyl group or pyrazolopyrimidinyl group; a sulfur-containing aromatic heterocyclic group such as thienyl group or benzothienyl group; and an oxygen-containing aromatic heterocyclic group such as furyl group, pyranyl group, cyclopentapyranyl group, benzofuryl group or isobenzofuryl group; an aromatic heterocyclic group containing two or more different hetero atoms, such as thiazolyl group, isothiazolyl group, benzothiazolyl group, benzothiadiazolyl group, phenothiazinyl group, isoxazolyl group, furazanyl group, phenoxazinyl group, oxazolyl group, isoxazolyl group, benzoxazolyl group, oxadiazolyl group, pyrazolooxazolyl group, imidazothiazolyl group, thienofuranyl group, furopyrrolyl group or pyridoxazinyl group, of which a preferred example is thienyl group, furyl group, pyridinyl group, pyridazinyl group, pyrimidinyl group or pyrazinyl group.

[0050] The "3-membered to 14-membered nitrogen-containing non-aromatic heterocyclic group" used in the specification of the present application means a monocyclic, bicyclic or tricyclic 3-membered to 14-membered non-aromatic heterocyclic group containing one or more nitrogen atoms. Preferable examples thereof include an azolidinyl group, azetizyl group, pyrrolidinyl group, pyrrolyl group, piperidinyl group, piperazinyl group, homopiperidinyl group, homopiperazinyl group, imidazolyl group, pyrazolidinyl group, imidazolidinyl, morpholinyl group, thiomorpholinyl group, imidazolinyl group, oxazolinyl group and quinuclidinyl group. The nitrogen-containing non-aromatic heterocyclic group also includes a group derived from a pyridone ring and a non-aromatic condensed ring (such as a group derived from a phthalimide ring or succinimide ring).

[0051] The " C_{2-22} alkanoyl group" used in the specification of the present application means a group corresponding to the above-defined " C_{1-22} alkyl group" in which the end thereof is a carbonyl group. Examples thereof include acetyl group, propionyl group, butyryl group, iso-butyryl group, valeryl group, iso-valeryl group, pivalyl group, caproyl group, decanoyl group, lauroyl group, myristoyl group, palmitoyl group, stearoyl group and arachidoyl group. Preferable examples thereof include alkanoyl groups having 2 to 6 carbon atoms such as acetyl group, propionyl group, butyryl group or iso-butyryl group.

[0052] The " C_{7-15} aroyl group" used in the specification of the present application means a group corresponding to the above-defined " C_{6-14} aryl group" or "5-membered to 14-membered heteroaryl group" to each of which end a carbonyl group is bonded. Examples thereof include benzoyl group, 1-naphthoyl group, 2-naphthoyl group, picolinoyl group, nicotinoyl group, isonicotinoyl group and furoyl group.

[0053] The " C_{3-23} unsaturated alkanoyl group" used in the specification of the present application means a group corresponding to the above-defined "unsaturated C_{2-22} alkyl group" to which end a carbonyl group is bonded. Examples thereof include an acryloyl group, propioloyl group, crotonoyl group, iso-crotonoyl group, oleoyl group and linolenoyl group. Preferable examples thereof include unsaturated alkanoyl groups having 2 to 6 carbon atoms and specifically an acryloyl group.

[0054] The " C_{7-22} aralkyl group" used in the specification of the present application means a group corresponding to the above-defined " C_{1-22} alkyl group" of which substitutable moiety is replaced by the above-defined " C_{6-14} aryl group" and being constituted of 7 to 22 carbon atoms. Specific examples thereof are benzyl group, phenethyl group, 3-phenylpropyl group, 4-phenylbutyl group, 1-naphthylmethyl group and 2-naphthylmethyl group, of which an aralkyl group having 7 to 10 carbon atoms such as benzyl group or phenethyl group is preferred.

[0055] The " C_{1-22} alkoxy group" used in the specification of the present application means a group corresponding to the above-defined " C_{1-22} alkyl group" to which end an oxygen atom is bonded. Suitable examples thereof are methoxy group, ethoxy group, n-propoxy group, iso-propoxy group, n-butoxy group, iso-butoxy group, sec-butoxy group, tert-butoxy group, n-pentyloxy group, iso-pentyloxy group, sec-pentyloxy group, n-hexyloxy group, iso-hexyloxy group, 1,1-dimethylpropoxy group, 1,2-dimethylpropoxy group, 2,2-dimethylpropoxy group, 2-ethylpropoxy group, 1-ethyl-2-methylpropoxy group, 1,1,2-trimethylpropoxy group, 1,2,2-trimethylpropoxy group, 1,1-dimethylbutoxy group, 1,2-dimethylbutoxy group, 2,2-dimethylbutoxy group, 2,3-dimethylbutoxy group, 1,3-dimethylbutoxy group, 2-ethylbutoxy group, 1,3-dimethylbutoxy group, 2-methylpentyloxy group, 3-methylpentyloxy group and hexyloxy group.

[0056] The "unsaturated C_{2-22} alkoxy group" used in the specification of the present application means a group corresponding to the above-defined "unsaturated C_{2-22} alkyl group" to which end an oxygen atom is bonded. Suitable examples thereof are vinyloxy group, allyloxy group, 1-propenyloxy group, isopropenyloxy group, 2-methyl-1-propenyloxy group, 2-methyl-2-propenyloxy group, 1-butenyloxy group, 2-butenyloxy group, 3-butenyloxy group, 1-pentyloxy group, 1-hexenyloxy group, 1,3-hexadienyloxy group, 1,5-hexadienyloxy group, propargyloxy group and 2-butyloxy group.

[0057] The " C_{6-14} aryloxy group" used in the specification of the present application means a group corresponding to the above-defined " C_{6-14} aryl group" to which end an oxygen atom is bonded. Specific examples thereof are phenoxy group, indenyloxy group, 1-naphthyloxy group, 2-naphthyloxy group, azulenyloxy group, heptalenyloxy group, indace-

nyloxy group, acenaphthyloxy group, fluorenyloxy group, phenalenyloxy group, phenanthrenyloxy group and anthracenyloxy group.

[0058] The "5-membered to 14-membered heteroaryloxy group" used in the specification of the present application means a group corresponding to the above-defined "5-membered to 14-membered heteroaryl group" to which end an oxygen atom is bonded. Specific examples thereof are pyrrolyloxy group, pyridinyloxy group, pyridazinyloxy group, pyrimidinyloxy group, pyrazinyloxy group, triazolyloxy group, tetrazolyloxy group, benzotriazolyloxy group, pyrazolyloxy group, imidazolyloxy group, benzimidazolyloxy group, indolyloxy group, isoindolyloxy group, indolizinyloxy group, purinyloxy group, indazolyloxy group, quinolinyloxy group, isoquinolinyloxy group, quinolizinyloxy group, phthalazinyloxy group, naphthyridinyloxy group, quinoxalinyloxy group, quinoxalinyloxy group, cinnolinyloxy group, pteridinyloxy group, imidazotriazinyloxy group, pyrazinopyridazinyloxy group, acridinyloxy group, phenanthridinyloxy group, carbazolyloxy group, carbazolyloxy group, perimidinyloxy group, phenanthrolinyloxy group, phenazinyloxy group, imidazopyridinyloxy group, imidazopyrimidinyloxy group, pyrazolopyridinyloxy group, pyrazolopyridinyloxy group, thienyloxy group, benzothienyloxy group, furyloxy group, pyranlyloxy group, cyclopentapyranlyloxy group, benzofuryloxy group, isobenzofuryloxy group, thiazolyloxy group, isothiazolyloxy group, benzothiazolyloxy group, benzothiadiazolyloxy group, phenothiazinyloxy group, isoxazolyloxy group, furazanyloxy group, phenoxazinyloxy group, oxazolyloxy group, isoxazoyloxy group, benzoxazolyloxy group, oxadiazolyloxy group, pyrazolooxazolyloxy group, imidazothiazolyloxy group, thienofuranyloxy group, furopyrrolyloxy group and pyridoxazinyloxy group, of which a preferred example is thienyloxy group, furyloxy group, pyridyloxy group, pyridazyloxy group, pyrimidyloxy group or pyrazolyloxy group.

[0059] The "5-membered to 14-membered heteroaryloxyalkyl group" used in the specification of the present application means a group corresponding to the above-defined "C₁₋₆ alkyl group" which is substituted with the above-defined "5-membered to 14-membered heteroaryloxy group".

[0060] The "C₁₋₂₂ alkylsulfonyl group" used in the specification of the present application means a sulfonyl group to which the above-defined "C₁₋₂₂ alkyl group" is bound. Specific examples thereof are methanesulfonyl group, ethanesulfonyl group, n-propanesulfonyl group and iso-propanesulfonyl group.

[0061] The "C₆₋₁₄ arylsulfonyl group" used in the specification of the present application means a sulfonyl group to which the above-defined "C₆₋₁₄ aryl group" is bound. Specific examples thereof are benzenesulfonyl group, 1-naphthalenesulfonyl group and 2-naphthalenesulfonyl group.

[0062] The "C₁₋₂₂ alkylsulfonyloxy group" used in the specification of the present application means a group corresponding to the above-defined "C₁₋₂₂ alkylsulfonyl group" to which end an oxygen atom is bonded. Examples thereof are methylsulfonyloxy group, ethylsulfonyloxy group, n-propylsulfonyloxy group and iso-propylsulfonyloxy group.

[0063] Examples of the substituent in the term "may have a substituent" used in the specification of the present application include those selected from the group consisting of:

- (1) halogen atom;
- (2) hydroxyl group;
- (3) thiol group;
- (4) nitro group;
- (5) nitroso group;
- (6) cyano group;
- (7) carboxyl group;
- (8) sulfonyloxy group;
- (9) amino group;
- (10) a C₁₋₂₂ alkyl group (for example, methyl group, ethyl group, n-propyl group, iso-propyl group, n-butyl group, iso-butyl group, sec-butyl group and tert-butyl group);
- (11) an unsaturated C₂₋₂₂ alkyl group (for example, vinyl group, allyl group, 1-propenyl group, isopropenyl group, ethynyl group, 1-propynyl group, 2-propynyl group, 1-butylnyl group, 2-butylnyl group and 3-butylnyl group);
- (12) a C₆₋₁₄ aryl group (for example, phenyl group, 1-naphthyl group and 2-naphthyl group);
- (13) a 5-membered to 14-membered heteroaryl group (for example, thienyl group, furyl group, pyridinyl group, pyridazinyl group, pyrimidinyl group and pyrazinyl group);
- (14) a 3-membered to 14-membered nitrogen-containing non-aromatic heterocyclic group (for example, aziridinyl group, azetidyl group, pyrrolidinyl group, pyrrolyl group, piperidinyl group, piperazinyl group, imidazolyl group, pyrazolidinyl group, imidazolidinyl, morpholinyl group, imidazoliny group, oxazoliny group and quinuclidinyl group);
- (15) a C₁₋₂₂ alkoxy group (for example, methoxy group, ethoxy group, n-propoxy group, iso-propoxy group, sec-propoxy group, n-butoxy group, iso-butoxy group, sec-butoxy group and tert-butoxy group);
- (16) a C₆₋₁₄ aryloxy group (for example, phenoxy group, 1-naphthyloxy group and 2-naphthyloxy group);
- (17) a C₇₋₂₂ aralkyloxy group (for example, benzyloxy group, phenethyloxy group, 3-phenylpropyloxy group, 4-phenylbutyloxy group, 1-naphthylmethyloxy group and 2-naphthylmethyloxy group);
- (18) a 5-membered to 14-membered heteroaryloxy group (for example, thienyloxy group, furyloxy group, pyridinyloxy

group, pyridazinyloxy group, pyrimidinyloxy group and pyrazinyloxy group);

(19) a C₂₋₂₃ alkanoyl group (for example, acetyl group, propionyl group, butyryl group, iso-butyryl group, valeryl group, iso-valeryl group, pivalyl group, caproyl group, decanoyl group, lauroyl group, myristoyl group, palmitoyl group, stearoyl group and arachidoyl group);

(20) a C₇₋₁₅ aroyl group (for example, benzoyl group, 1-naphthoyl group and 2-naphthoyl group);

(21) a C₃₋₂₃ unsaturated alkanoyl group (for example, acryloyl group, propioloyl group, crotonoyl group, iso-crotonoyl group, oleoyl group and linolenoyl group);

(22) a C₂₋₂₃ alkanoyloxy group (for example, acetoxyl group, propionyloxy group and acryloxy group);

(23) a C₂₋₂₂ alkoxy carbonyl group (for example, methoxycarbonyl group, ethoxycarbonyl group, n-propoxycarbonyl group, iso-propoxycarbonyl group, n-butoxycarbonyl group, iso-butoxycarbonyl group, sec-butoxycarbonyl group and tert-butoxycarbonyl group);

(24) an unsaturated C₃₋₂₂ alkoxy carbonyl group (for example, vinyloxy carbonyl group, aryloxy carbonyl group, 1-propenyloxy carbonyl group, isopropenyloxy carbonyl group, propargyloxy carbonyl group and 2-butyloxy carbonyl group);

(25) a C₁₋₂₂ alkylsulfonyl group (for example, methanesulfonyl group, ethanesulfonyl group, n-propanesulfonyl group and iso-propanesulfonyl group);

(26) a C₆₋₁₄ arylsulfonyl group (for example, benzenesulfonyl group, 1-naphthalenesulfonyl group and 2-naphthalenesulfonyl group); and

(27) a C₁₋₂₂ alkylsulfonyloxy group (for example, methanesulfonyloxy group, ethanesulfonyloxy group, n-propanesulfonyloxy group and iso-propanesulfonyloxy group).

Examples

Reference Example 1 Production of starting material, a macrolide compound 11107B

[0064] One loopful of the slant culture (ISP-2 medium) of *Streptomyces* sp. Mer-11107 strain (FERM BP-7812) was inoculated into a 500 mL Erlenmeyer flask containing 50 mL of seed medium (2% of glucose, 1% of ESUSAN-MEAT manufactured by Ajinomoto Co. Ltd., 0.5% of yeast extract (manufactured by Oriental Yeast Co., Ltd.), 0.25% of sodium chloride, 0.32% of calcium carbonate, pH 6.8 before sterilization), and it was incubated at 28°C for two days to give the first seed culture broth. 0.1 mL of the culture broth was inoculated into a 500 mL Erlenmeyer flask containing 100 mL of the same seed medium and it was incubated at 28°C for one day to give the second seed culture broth. The second seed culture broth (800 mL) thus obtained was inoculated into a 200 L tank containing 100 L of a production medium (5% of soluble starch, 0.8% of Pharmamedia, 0.8% of gluten meal, 0.5% of yeast extract and 0.1% of calcium carbonate, pH 6.8 before sterilized) and it was cultured for five days with flowing air and stirring under the conditions of a culture temperature of 28°C, an agitation rotation of 90 rpm, a quantity of aeration of 1.0 vvm and an internal pressure of 20 kPa, to give a culture broth.

[0065] A part of the culture broth (10L) thus obtained was extracted with 10 L of 1-butanol, and then the resulting butanol layer was evaporated to dryness, to give 100 g of crude active fraction. The crude active fraction was applied on Sephadex LH-20 (1500 mL; manufactured by Pharmacia Co. Ltd.), and eluted with tetrahydrofuran-methanol (1:1) as a solvent. An eluted fraction from 540 mL to 660 mL was concentrated to dryness, to give a residue (660 mg). The resulting residue was dissolved in a mixture of ethyl acetate and methanol (9:1; v/v) and subjected to silica gel column chromatography (WAKO GEL C-200, 50 g). The column was eluted with a mixture (2 L) consisting of n-hexane and ethyl acetate (1:9, v/v), the fractions eluted from 468 mL to 1260 mL were collected, evaporated to give 25 mg of a crude active fraction.

[0066] The obtained crude active fraction was subjected to preparative high performance liquid chromatography (HPLC) under the following preparative HPLC condition (A), and the fractions eluted at the retention time of 34 minutes were collected. After removing acetonitrile, the respective fractions were desalted by HPLC under the following preparative HPLC condition (B) to give the macrolide compound 11107B (Retention time: 37 minutes, 6 mg).

Preparative HPLC conditions A:

[0067]

Column: YMC-PACK ODS-AM SH-343-5ÅM, φ20 mm × 250 mm (manufactured by YMC Co.)
 Temperature: room temperature
 Flow rate: 10 mL/min.
 Detection: 240 nm
 Eluent: acetonitrile/0.15% aqueous potassium dihydrogenphosphate (pH 3.5) (2:8 to 8:2, v/v, 0 to 50 min.,

linear gradient)

Preparative HPLC conditions B:

[0068]

Column: YMC-PACKODS-AMSH-343-5AM, ϕ 20mm \times 250mm (manufactured by YMC Co.)
 Temperature: room temperature
 Flow rate: 10 ml/min.
 Detection: 240 nm
 Eluent: methanol/water (2: 8 to 10: 0, v/v, 0 to 40 minutes, linear gradient)

Example 1 Determination of the nucleotide sequence of a gene derived from *Streptomyces* sp. A-1544 strain (FERM BP-8446) (1) Preparation of a DNA of *Streptomyces* sp. A-1544 strain chromosome

[0069] The A-1544 strain was inoculated into a medium containing 1% of glucose, 0.4% of malt extract and 1% of yeast extract and incubated at 28°C for 3 days. The obtained culture broth was centrifuged at 3000 rpm for 10 minutes to collect the mycelia. A chromosome DNA was prepared using Blood & Cell Culture kit (QIAGEN Co.) from the mycelia.

(2) Cloning of a partial sequence of a DNA encoding a protein having the activity in hydroxylating the 16-position of the macrolide compound 11107

[0070] Mix primers 5Dm-3F (sequence no. 4) and 5Dm-3R (sequence No. 5) were designed and produced on reference to the amino acid sequence assumed to be that of the cytochrome P450 (CYP105D5) of *Streptomyces coelicolor* A3 (2).

[0071] In order to promote reactivity taking the fluctuation of a codon into account, mixed bases S (=C+G) and Y (=C+T) were used.

[0072] Next, these two types of primers (5Dm-3F and 5Dm-3R) and the A-1544 strain chromosome DNA obtained in the above (1) as a template, were used to run a PCR reaction. The PCR reaction was accomplished by repeating a three-stage reaction including denaturing run at 98°C for 20 seconds, annealing run at 50°C for 2 minutes and extension run at 68°C for 30 seconds 35 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and a PCR amplifier (T Gradient, Biometra Co.). As a result, a DNA fragment (hereinafter referred to as a DNA fragment-A1) having a size of about 500 bp was amplified. It is highly possible that this DNA fragment-A1 is a part of the DNA encoding a protein having hydroxylating activity. The DNA fragment-A1 amplified by a PCR reaction was recovered from the reaction solution by SUPREC PCR (TAKARA HOLDINGS INC.).

[0073] In order to obtain the DNA fragment-A1 in an amount enough to analyze the nucleotide sequence of the obtained DNA fragment-A1, the DNA fragment was combined with a plasmid vector pT7Blue T (Novagen Co.) by using DNA Ligation kit ver.2 (TAKARA HOLDINGS INC.) to transform *E. coli* JM109 strain. Thereafter, the transformed *E. coli* was selected using a L-broth agar media (1.0% bactotripton, 0.5% yeast extract, 0.5% NaCl, 1.5% agar) containing ampicillin (50 μ g/mL), X-gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside; 40 μ g/mL) and IPTG (isopropyl- β -D-thiogalactopyranoside; 100 μ M). The colony of the transformed *E. coli* thus isolated was cultured in a L-broth liquid medium (1% bactotripton, 0.5% yeast extract, 0.5% NaCl) containing ampicillin (50 μ g/mL). A plasmid DNA was separated from the mycelia of the proliferated transformed *E. coli* and purified by using a plasmid purifying kit (QIA filter Plasmid Midi Kit, QIAGEN Co.), to obtain enough amount of the DNA fragment-A1.

(3) Analysis of the nucleotide sequence of the cloned DNA fragment-A1

[0074] The nucleotide sequence of the DNA fragment-A1 obtained in the above (2) was analyzed using a DNA nucleotide sequence analyzer (PE Biosystems 377XL) according to a dye terminator cycle sequence method. As the result of the nucleotide sequence analysis, it was clarified that the DNA fragment-A1 amplified by a PCR reaction had an exact size of 528 bp though it had been found to have a size of about 500 bp by the above measurement using electrophoresis (see the nucleotide sequence 1775 to nucleotide sequence 2302 of the sequence No. 1). Since DNA sequences corresponding to the two types of primers used in the above PCR reaction were found at both ends of the above cloned 528 bp DNA sequence, it was clarified that the DNA fragment-A1 was singularly amplified by these two types of primers (5Dm-3F and 5Dm-3R) in the above PCR reaction.

(4) Analysis of the neighboring region of the DNA fragment-A1

[0075] As mentioned above, the partial sequence of the DNA encoding a protein which was derived from the A-1544

strain and had hydroxylating activity. Therefore, the amplification, cloning and sequence analysis of the nucleotide sequence in the neighboring region extending from the upstream side to downstream side of the cloned fragment were accomplished by an inverse PCR method (Cell Technology vol. 14, p.591-593, 1995). Specifically, the A-1544 strain chromosome DNA (see the above (1)) was digested by respective restriction enzymes PstI and Sall in a H buffer solution (50 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 10 mM dithiothreitol and 100 mM NaCl). The obtained each DNA fragment cut by the restriction enzymes was self-circularized by using DNA Ligation Kit ver.2 (TAKARA HOLDINGS INC.).

[0076] On the other hand, primers (6PIN-2F (sequence No. 6) and 6PIN-2R (sequence No. 7) were designed and produced based on the nucleotide sequence of the DNA fragment-A1.

[0077] Next, these two primers (6PIN-2F and 6PIN-2R) and the above self-cyclized A-1599 strain chromosome DNA as a template, were used to run a PCR reaction. In the PCR reaction, the cycle of a two-stage reaction involving denaturing run at 98°C for 20 seconds and annealing and extension run at 68°C for 5 minutes was repeated 35 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and a PCR amplifier (T Gradient, Biometra Co.).

[0078] As a result, a DNA fragment (DNA fragment-B1) about 3.5 kbp in size and a DNA fragment (DNA fragment-C1) about 2.8 kbp in size were amplified. It was highly possible that these DNA fragments were a DNA encoding a protein having hydroxylating activity and a DNA having a DNA sequence including the upstream and downstream regions of the former DNA.

[0079] The DNA fragment-B1 and the DNA fragment-C1 were recovered from the PCR amplified reaction solution by SUPREC PCR (TAKARA HOLDINGS INC.). Next, as to the obtained DNA fragment-B1 and DNA fragment-C1, in order to obtain each DNA fragment in an amount enough to analyze the nucleotide sequence of the obtained DNA fragment, a plasmid vector pT7Blue T (Novagen Co.), DNA Ligation kit ver.2 (TAKARA HOLDINGS INC.), E. coli JM109 strain and a plasmid purifying kit (QIA filter Plasmid Midi kit, QIAGEN Co.) were used in the same manner as the above (2), to obtain enough amount of each DNA fragment.

(5) Analysis of each nucleotide sequence of the DNA fragment-B1 (about 3.5 kbp in size) and the DNA fragment-C1 (about 2.8 kbp in size)

[0080] Each nucleotide sequence of the DNA fragment-B1 and DNA fragment-C1 obtained in the above (4) was analyzed using a DNA nucleotide sequence analyzer (PE Biosystems 377XL) according to a dye terminator cycle sequence method. The nucleotide sequence was thus analyzed to obtain the information of the nucleotide sequence of 3793 bp shown in the sequence No. 1 from each sequence of the DNA fragment-B1 and DNA fragment-C1.

[0081] An open reading frame (ORF) in this 3793 bp was retrieved, to find that the two proteins were coded. Each amino acid sequence of these proteins was retrieved by the BLAST search, and as a result, an ORF (hereinafter referred to as psmA) coding a protein consisting of 409 amino acids having high homology to cytochrome P450 existed in the base 1322 to base 2548 of the sequence No. 1. The psmA had the highest homology (homology: 72.6%) to the amino acid sequence assumed to be that of cytochrome P450 (CYP105D5) of the *Streptomyces coelicolor* A3 (2) and to the amino acid sequence assumed to be that of cytochrome P450 (CYP105D4) of the *Streptomyces lividans*, and also had a relatively high homology (homology: 69.4%) to cytochrome P450 soy (Soy C) of *Streptomyces griseus*. It was considered from this fact that the psmA was highly possibly a gene coding hydroxylating enzyme of the cytochrome P-450 type.

[0082] Also, an ORF (hereinafter referred to as psmB) encoding a protein having a high homology to ferredoxin of a 3F-4S type existed just downstream (the base 2564 to base 2761 of the sequence No. 1) of the psmA. The protein encoded by the psmB consists of 66 amino acids, and had the highest homology (83.3%) to the amino acid sequence assumed to be that of ferredoxin just downstream of the amino acid sequence assumed to be that of cytochrome P450 (CYP105D5) of the *Streptomyces coelicolor* A3 (2) and a relatively higher homology (homology: 57.6%) to ferredoxin soy (soyB) of *Streptomyces griseus*. Therefore, it was considered that the psmB serves to transfer electrons and codes ferredoxin participating in hydroxylation together with the psmA. Example 2 Production of a transformant having the psmA and the psmB

(1) Preparation of a DNA fragment containing both the psmA and the psmB derived from the A-1544 strain

[0083] A primer DM-NdeF (sequence No. 8) obtained by adding a NdeI site to the 5' terminal and a primer DM-SpeR (sequence No. 9) obtained by adding a SpeI site to the 5' terminal were designed and produced on reference to the nucleotide sequence of the sequence No. 1 analyzed in Example 1. Next, these two types of primers (DM-NdeF and DM-SpeR) and the A-1544 strain chromosome DNA obtained in Example 1(1) as a template, were used to run a PCR reaction. The PCR reaction was accomplished by repeating a two-stage reaction including denaturing carried out at 98°C for 20 seconds and annealing and elongation carried out at 68°C for 2 minutes 30 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and a PCR amplifier (T Gradient, Biometra Co.).

[0084] As a result, a DNA fragment (hereinafter referred to as a DNA fragment-D1) having a size of about 1.5 kbp and containing the psmA and the psmB was amplified. The DNA fragment-D1 was recovered from this PCR amplified

reaction solution by SUPREC PCR (TAKARA HOLDINGS INC.).

(2) Architecture of a plasmid pTC-DM



5 **[0085]** pT7NS-CamAB (see WO03/087381) was digested by respective restriction enzymes NdeI and SpeI in a H
buffer solution (50 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 10 mM dithiothreitol and 100 mM NaCl) to obtain a plasmid
digested products. Similarly, the DNA fragment-D1 obtained in the above (1) was digested by respective restriction
enzymes NdeI and SpeI. The obtained digested product of the DNA fragment-D1 and the plasmid digested product were
coupled using DNA Ligation Kit ver.2 (TAKARA HOLDINGS INC.). This resulted in the formation of a plasmid (referred
10 to as a plasmid pTC-DM) about 9.5 kbp in size which was an architecture of a combination of the DNA fragment-D1
containing both the psmA and the psmB therein and the plasmid pT7NS-CamAB.

(3) Preparation of *E. coli* transforming strain BL21 (DE3) /pTC-DM

15 **[0086]** Using the plasmid pTC-DM prepared in the above (2), a competent cell (Novagen) of *Coli* bacillus BL21 (DE3) was transformed. *E. coli* BL21 (DE3) /pTC-DM strain transformed by the plasmid pTC-DM was obtained.

Example 3: Conversion of ME-265 into ME-282 represented by the following formulae by the *E. coli* transformant having the *psmA* and the *psmB*

[0087]



(1) Preparation of a transformant reaction solution

[0088] The transformed *E. coli* BL21 (DE3) /pTC-DM strain obtained in Example 2(3) and a frozen seed of a BL21 (DE3)/pT7NS-CamAB strain were inoculated into a 15 mL test tube containing 3 mL of a L-broth medium (1.0% bactotripton, 0.5% yeast extract, 0.5% NaCl) containing 50 µg/mL of ampicillin and shake-cultured at 37°C for 20 hours. 500

μL of the seed culture broth was inoculated into a 250 mL Erlenmeyer flask containing 50 mL of a L-broth medium (1.0% bactotripton, 0.5% yeast extract, 0.5% NaCl) containing 50 μg/mL of ampicillin and shake-cultured at 32°C for 3 hours. Then, 50 μL of 100 mM IPTG (isopropyl-β-D-thiogalactopyranoside) and 50 μL of 80 mg/mL 5-aminolevulinic acid were successively added thereto, and the medium was shake-cultured at 32°C for 6 hours. The obtained culture broth was centrifuged (5000 rpm, 10 minutes) to collect the mycelia. The mycelia were then suspended in 1.75 mL of a 100 mM phosphate buffer solution (pH 6.1), and 250 μL of 80% glycerol and 50 μL of 8 mg/mL ME-265 were added thereto. The conversion reaction solution thus obtained was reacted at 28°C for 24 hours. 200 μL of the reaction solution was extracted with 1 mL of acetonitrile and the extract was subjected to HPLC to measure each amount of ME-265 and ME-282. The results are shown in Table 3.

[0089] Also, the details of the condition of HPLC are shown below.

Analyzer: Shimadzu HPLC 10Avp

Column: CAPCELL PAK C18 SG120 (φ4.6 mm x250 mm)

Mobile phase: 45% acetonitrile (0 to 15 minutes)

60% acetonitrile (15 to 30 minutes)

45% acetonitrile (30 to 45 minutes)

Flow rate: 1 mL/min.

Detection: UV 240 nm

Injection capacity: 10 μL

Column temperature: 40°C

Analyzing time: 45 minutes

Retention time: ME-265 24.8 minutes

ME-282 12.7 minutes

Table 3

mg/L	BL21(DE3)/pT7NS-CamAB	BL21(DE3)/pTC-DM
ME-265	143	0
ME-282	0	130

(2) Isolation of ME-282 from the transformant reaction solution

[0090] 4 mL of water was added to 1.8 mL of the reaction solution that had been reacted for 24 hours and the reaction solution was then extracted with 8 mL of ethyl acetate once and with 4 mL of ethyl acetate twice. The ethyl acetate layers were combined, dried over anhydrous sodium sulfate and then the solvent was removed. The resulting residue was purified by thin layer chromatography (MERCK Silicagel 60 F254.0.25 mm, developing solution: hexane: ethyl acetate=1:2), to give 0.2 mg of ME-282.

[0091] ¹H-NMR spectrum (CD₃OD, 500MHz): δ ppm (integral, multiplicity, coupling constant J(Hz)):

0.87(3H,d,J=7.0Hz), 0.90(3H,d,J=7.0Hz), 0.94(3H,t,J=7.3Hz), 0.97(3H,d,J=6.6Hz), 1.21-1.26(1H,m), 1.29-1.37(3H,m), 1.39(3H,s), 1.94-1.52(2H,m), 1.60-1.64(1H,m), 1.65(1H,dd,J=6.2,13.9Hz), 1.77(3H,d,J=1.1Hz), 1.86(1H,dd,J=5.4,13.9Hz), 1.89-1.99(1H,m), 2.00(3H,s), 2.43(1H,dd,J=5.5,13.9Hz), 2.50-2.60(1H,m), 2.56(1H,dd,J=3.3,13.9Hz), 2.66(1H,dd,J=2.2,7.7Hz), 2.89(1H,dt,J=2.2,6.2Hz), 3.52(1H,dt,J=4.8,8.4Hz), 3.75-3.80(1H,m), 4.90(1H,overlapped with-D₂O), 5.01(1H,d,J=10.6Hz), 5.42(1H,dd,J=9.2,15.0Hz), 6.13(1H,d,J=10.6Hz), 6.52(1H,dd,J=11.0,15.0Hz).

[0092] As a result, a peak estimated as that of ME-282 was not observed in the case of the E. coli BL21 (DE3) /pT7NS-CamAB strain used as a control, whereas ME-265 was almost consumed and a peak estimated as that of ME-282 was obtained in the case of the BL21 (DE3) /pTC-DM strain containing the psmA and psmB. This fact suggests that the psmA and the psmB participate in the conversion of ME-265 into ME-282.

Example 4: Conversion of the macrolide compound 11107B into the macrolide compound 11107D by the E. coli transformant having the psmA and the psmB

(1) Preparation of a transformant reaction solution

[0093] A test using the macrolide compound 11107B as a substrate was made in the same manner as Example 3. The transformed E. coli BL21(DE3)/pTC-DM strain obtained in Example 2(3) and a frozen seed of a BL21 (DE3) /pT7NS-CamAB strain were inoculated into a 15 mL test tube containing 3 mL of a L-broth medium (1.0% bactotripton, 0.5% yeast extract, 0.5%NaCl) containing 50 μg/mL of ampicillin and shake-cultured at 30°C for 20 hours. 500 μL of the seed culture broth was inoculated into a 250 mL Erlenmeyer flask containing 50 mL of a L-broth medium (1.0% bactotripton,

0.5% yeast extract, 0.5% NaCl) containing 50 µg/mL of ampicillin and shake-cultured at 28°C for 5 hours. Then, 50 µL of 100 mM IPTG (isopropyl-β-D-thiogalactopyranoside) and 50 µL of 80mg/mL 5-aminolevulinic acid were successively added and the medium was shake-cultured at 25°C for 20 hours. The obtained culture broth was centrifuged (5000 rpm, 10 minutes) to collect the mycelia. The mycelia were then suspended in 1.75 mL of a 100 mM phosphate buffer solution (pH 6.1), and 250 µL of 80% glycerol and 50 µL of 40 mg/mL 11107B were added thereto. The conversion reaction solution thus obtained was reacted at 28°C for 24 hours. 200 µL of the reaction solution was extracted with 1 mL of acetonitrile and the extract was subjected to HPLC to measure each amount of the macrolide compound 11107B and the macrolide compound 11107D. The results are shown in Table 4. Also, the details of the condition of HPLC are shown below.

Analyzer: Shimadzu HPLC 10Avp

Column: CAPCELL PAK C18 SG120 (φ4.6 mm × 250 mm)

Mobile phase: 35% acetonitrile (0 to 10 minutes)

35% to 65% acetonitrile (10 to 12 minutes)

65% acetonitrile (12 to 15 minutes)

35% acetonitrile (15 to 20 minutes)

Flow rate: 1 mL/min.

Detection: UV 240 nm

Injection capacity: 10 µL

Column temperature: 40°C

Analyzing time: 20 minutes

Retention time: 11107B 14.3 minutes

11107D 7.9 minutes

Table 4

mg/L	BL21(DE3)/pT7NS-CamAB	BL21(DE3)/pTC-DM
11107B	636	619
11107D	0	71

(2) Isolation of the macrolide compound 11107D from the transformant reaction solution

[0094] 4 mL of water was added to 1.8 mL of the reaction solution that had been reacted for 24 hours and the mixture was then extracted with 8 mL of ethyl acetate once and with 4 mL of ethyl acetate twice. The ethyl acetate layers were combined, dried over anhydrous sodium sulfate and the solvent was removed. The resulting residue was purified by thin layer chromatography (MERCK Silicagel 60 F254 0.25 mm, developing solution: ethyl acetate) to obtain 0.1 mg of 11107D.

[0095] ¹H-NMR spectrum (CD₃OD, 500MHz) : δ ppm (integral, multiplicity, coupling constant J(Hz)):

0.87(3H,d,J=7.0Hz), 0.88(3H,d,J=7.0Hz), 0.93(3H,t,J=7.0Hz), 1.18(3H,s), 1.18-1.69(8H,m), 1.33(3H,s), 1.77(3H,d,J=1.1Hz), 1.82-1.90(1H,m), 2.05(3H,s), 2.49-2.60(3H,m), 2.66(1H,dd,J=2.2,8.2Hz), 2.89(1H,dt,J=2.4,5.7Hz), 3.52(1H,dt,J=4.8,8.3Hz), 3.73-3.82(1H,m), 5.09(1H,d,J=9.8Hz), 5.05(1H,d,J=10.6Hz), 5.56(1H,dd,J=9.8,15.2Hz), 5.70(1H,dd,J=9.8,15.2Hz), 5.86(1H,d,J=15.2Hz), 6.3(1H,d,J=10.8Hz), 6.52(1H,dd,J=10.8,15.2Hz).

[0096] As a result, a peak estimated as that of the macrolide compound 11107D was not observed in the case of the E. coli BL21 (DE3)/pT7NS-CamAB strain used as a control, whereas a peak estimated as that of the macrolide compound 11107D was obtained in the case of the BL21 (DE3)/pTC-DM strain containing the psmA and psmB. This fact suggests that the psmA and the psmB participate in the conversion of macrolide compound 11107B into the macrolide compound 11107D.

Example 5: Conversion test using an A-1544 self-cloning strain

(1) Preparation of a DNA fragment containing both the psmA and the psmB derived from the A-1544 strain

[0097] A primer DM-BglIF (sequence No. 10) obtained by adding a BglII site to the 5' terminal and a primer DM-BglIR (sequence No. 11) obtained by adding a BglII site to the 5' terminal were designed and produced on reference to the nucleotide sequence of the sequence No. 1 analyzed in Example 1.

[0098] Next, these two types of primers (DM-BglIF and DM-BglIR) and the A-1544 strain chromosome DNA obtained in Example 1(1) as a template, were used to run a PCR reaction. The PCR reaction was accomplished by repeating a three-stage reaction including denaturing carried out at 98°C for 20 seconds, annealing carried out at 63°C for 30 seconds

and elongation carried out at 68°C for 4 minutes 30 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and a PCR amplifier (T Gradient, Biometra Co.).

[0099] As a result, a DNA fragment (hereinafter referred to as a DNA fragment-E1) having a size of about 3.5 kbp and containing the psmA and the psmB was amplified. This PCR amplified reaction solution was subjected to agarose gel electrophoresis to fractionate. The above DNA fragment-E1 about 3.5 kbp in size was cut out of the agarose gel and recovered by SUPREC 01 (TAKARA HOLDINGS INC.).

(2) Architecture of a plasmid pIJDMG

[0100] pIJ702 was digested by a restriction enzyme BglII in a H buffer solution (50 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 10 mM dithiothreitol and 100 mM NaCl) to obtain a plasmid digested products. Similarly, the DNA fragment-E1 obtained in the above (1) was digested by a restriction enzyme BglII. The obtained digested product of the DNA fragment-E1 and the plasmid-digested product were bound using DNA Ligation Kit ver.2 (TAKARA HOLDINGS INC.). The resulted in the formation of a plasmid (referred to as a plasmid pIJDMG) about 8.5 kbp in size which was an architecture of a combination of the DNA fragment-E1 containing both the psmA and the psmB therein and the plasmid pIJ702.

(3) Preparation of a self-cloning strain A-1544/pIJDMG strain

[0101] Using the plasmid pIJDMG prepared in the above (2), an A-1544 strain was transformed according to the method described in Genetic Manipulation of Streptomyces: A Laboratory Manual. John Innes Foundation, Norwich, 1985. An A-1544/pIJDMG strain was thus obtained by transformation using the plasmid pIJDMG.

Example 6: Conversion of 11107B into 11107D by a self-cloning strain

[0102] The transformed A-1544/pIJDMG strain obtained in Example 5(3), A-1544/pIJ702 strain and a frozen seed of the original A-1544 strain were inoculated into a 250 mL Erlenmeyer flask containing 50 mL of a SMN medium (Stabilose 2%, glucose 2%, ESUSAN-MEAT 2%, yeast extract 0.5%, NaCl 0.25%, CaCO₃ 0.32%, pH 7.4) containing 25 µg/mL of thiostrepton, and shake-cultured at 28°C for 48 hours (seed culture, no addition of thiostrepton to the A-1544 strain). 0.5 mL of the obtained seed culture broth was inoculated into a 250 mL Erlenmeyer flask containing 50 mL of a SMN medium containing 25 µg/mL of thiostrepton and shake-cultured at 28°C for 72 hours (no addition of thiostrepton to the A-1544 strain). The obtained culture broth was dispensed in 2 mL portions, and 100 µL of a 1M phosphate buffer solution (pH 6.5) and 50 µL of 40 mg/mL of 11107B were added thereto. The conversion culture broth thus obtained was reacted at 28°C for 12 hours. 200 µL of the reaction solution was extracted with 1 mL of acetonitrile and the extract was subjected to HPLC to measure each amount of 11107B and 11107D. The results are shown in Table 5. Also, the details of the condition of HPLC are shown below.

Analyzer: Shimadzu HPLC 10Avp

Column: CAPCELL PAK C18 SG120 (φ4.6 mm x250 mm)

Mobile phase: 35% acetonitrile (0 to 10 minutes)

35% to 65% acetonitrile (10 to 12 minutes)

65% acetonitrile (12 to 15 minutes)

35% acetonitrile (15 to 20 minutes)

Flow rate: 1 mL/min.

Detection: UV 240 nm

Injection capacity: 10 µL

Column temperature: 40°C

Analyzing time: 20 minutes

Retention time: 11107B 14.3 minutes

11107D 7.9 minutes

Table 5

mg/L	A-1544 strain	A-1544/pIJ702 strain	A-1544/pIJDMG strain
11107B	496	651	14
11107D	196	0	535

[0103] As a result, the A-1544/pIJDMG strain obtained by transformation of the plasmid containing the psmA and the psmB exhibited conversion activity about 2.7 times that of the original A-1544 strain by a reaction run for 12 hours. This

fact suggests that the self-cloning of the psmA and psmB contributes to the conversion of the macrolide compound 11107B into the macrolide compound 11107D.

Example 7 Determination of the nucleotide sequence of a gene derived from *Streptomyces* sp. Mer-11107 strain (FERM BP-7812)

(1) Preparation of a DNA of *Streptomyces* sp. Mer-11107 strain chromosome

[0104] The Mer-11107 strain was inoculated into a medium containing 1% of glucose, 0.4% of malt extract and 1% of yeast extract and cultured at 28°C for 3 days. The obtained culture broth was centrifuged at 3000 rpm for 10 minutes to collect the mycelia. A chromosome DNA was prepared using Blood & Cell Culture kit (QIAGEN Co.) from the mycelia.

(2) Cloning of a partial sequence of a DNA encoding a protein having the activity in hydroxylating the 16-position of the macrolide compound 11107

[0105] Mix primers 5Dm-3F (sequence no. 4) and 5D-1R (sequence No. 12) were designed and produced on reference to the amino acid sequence estimated as that of the cytochrome P450 (CYP105D5) of *Streptomyces coelicolor* A3(2).

[0106] In order to promote reactivity taking the fluctuation of a codon into account, mixed bases S (=C+G) and Y (=C+T) were used.

[0107] Next, these two types of primers (5Dm-3F and 5D-1R) and the Mer-11107 strain chromosome DNA obtained in the above (1) as a template, were used to run a PCR reaction. The PCR reaction was accomplished by repeating a three-stage reaction including denaturing carried out at 98°C for 20 seconds, annealing carried out at 50°C for 2 minutes and elongation carried out at 68°C for 30 seconds 35 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and a PCR amplifier (T Gradient, Biometra Co.). As a result, a DNA fragment (hereinafter referred to as a DNA fragment-A2) having a size of about 300 bp was amplified. It is highly possible that the DNA fragment-A2 is a part of the DNA encoding a protein having hydroxylating activity. The DNA fragment-A2 amplified by a PCR reaction was recovered from the reaction solution by SUPREC PCR (TAKARA HOLDINGS INC.).

[0108] In order to obtain the DNA fragment-A2 in an amount enough to analyze the nucleotide sequence of the obtained DNA fragment-A2, the DNA fragment was bound with a plasmid vector pT7Blue T (Novagen Co.) by using DNA Ligation kit ver.2 (TAKARA HOLDINGS INC.) to transform *E. coli* JM109 strain. Thereafter, the transformed *E. coli* was selected using a L-broth agar media (1.0% bactotripton, 0.5% yeast extract, 0.5% NaCl, 1.5% agar) containing ampicillin (50 µg/mL), X-gal (5-bromo-4-chloro-3-indolyl-β-D-galactoside; 40 µg/mL) and IPTG (isopropyl-β-D-thiogalactopyranoside; 100 µM). The colony of the transformed *E. coli* thus isolated was cultured in a L-broth liquid medium (1% bactotripton, 0.5% yeast extract, 0.5% NaCl) containing ampicillin (50 µg/mL). A plasmid DNA was separated from the mycelia of the proliferated transformed *E. coli* and purified by using a plasmid purifying kit (QIA filter Plasmid Midi Kit, QIAGEN Co.), to obtain enough amount of the DNA fragment-A2.

(3) Analysis of the nucleotide sequence of the cloned DNA fragment-A2

[0109] The nucleotide sequence of the DNA fragment-A2 obtained in the above (2) was analyzed using a DNA nucleotide sequence analyzer (PE Biosystems 377XL) according to a dye terminator cycle sequence method. As the result of the nucleotide sequence analysis, it was clarified that the DNA fragment-A2 amplified by a PCR reaction had an exact size of 325 bp though it had been found to have a size of about 300 bp by the above measurement using electrophoresis (see the nucleotide sequence 837 to nucleotide sequence 1161 of the sequence No. 2). Since DNA sequences corresponding to the two types of primers used in the above PCR reaction were found at both ends of the above cloned 325 bp DNA sequence, it was clarified that the DNA fragment-A2 was specifically amplified by these two types of primers (5Dm-3F and 5D-1R) in the above PCR reaction.

(4) Analysis of the neighboring region of the DNA fragment-A2

[0110] As mentioned above, the partial sequence of the DNA encoding a protein having the hydroxylating activity derived from the Mer-11107 strain was determined. Therefore, the amplification, cloning and sequence analysis of the nucleotide sequence in the neighboring region extending from the upstream side to downstream side of the cloned fragment were accomplished by an inverse PCR method (Cell Technology vol. 14, p.591-593, 1995).

Specifically, the Mer-11107 strain chromosome DNA (see the above (1)) was digested by a restriction enzyme BamHI in a K buffer solution (50 mM Tris-HCl, pH 8.5, 10 mM MgCl₂, 1 mM dithiothreitol and 100 mM KCl) and by a restriction enzyme Sall in a H buffer solution (50 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 1 mM dithiothreitol and 100 mM NaCl) respectively. The obtained each DNA fragment cut by the restriction enzymes was self-circularized using DNA Ligation

Kit ver.2 (TAKARA HOLDINGS INC.).

[0111] On the other hand, primers (7PIN-2F (sequence No. 13) and 6PIN-2R (sequence No. 7) were designed and produced based on the nucleotide sequence of the DNA fragment-A2.

[0112] Next, these two primers (7PIN-2F and 6PIN-2R) and the above self-circularized Mer-11107 strain chromosome DNA as a template, were used to run a PCR reaction. In the PCR reaction, the cycle of a two-stage reaction involving denaturing carried out at 98°C for 20 seconds and annealing and elongation carried out at 68°C for 5 minutes was repeated 35 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and a PCR amplifier (T Gradient, Biometra Co.).

[0113] As a result, a DNA fragment (DNA fragment-B2) about 1.3 kbp in size and a DNA fragment (DNA fragment-C2) about 1.4 kbp in size were amplified. It was highly possible that these DNA fragments were respectively a DNA encoding a protein having hydroxylating activity and a DNA having a DNA sequence including those in the upstream and downstream regions of the former DNA.

[0114] The DNA fragment-B2 and the DNA fragment-C2 were recovered from the PCR amplified reaction solution by SUPREC PCR (TAKARA HOLDINGS INC.). Next, as to the obtained DNA fragment-B2 and DNA fragment-C2, in order to obtain each DNA fragment in an amount enough to analyze the nucleotide sequence of the obtained DNA fragment, a plasmid vector pT7Blue T (Novagen Co.), DNA Ligation kit ver.2 (TAKARA HOLDINGS INC.), E. coli JM109 strain and a plasmid purifying kit (QIA filter Plasmid Midi Kit, QIAGEN Co.) were used in the same manner as the above (2), to obtain enough amount of each DNA fragment.

(5) Analysis of each nucleotide sequence of the DNA fragment-B2 (about 1.3 kbp in size) and the DNA fragment-C2 (about 1.4 kbp in size)

[0115] Each nucleotide sequence of the DNA fragment-B2 and DNA fragment-C2 obtained in the above (4) was analyzed using a DNA nucleotide sequence analyzer (PE Biosystems 377XL) according to a dye terminator cycle sequence method. The nucleotide sequence was thus analyzed to obtain the information of the nucleotide sequence of 2329 bp shown in the sequence No. 2 from each sequence of the DNA fragment-B2 and DNA fragment-C2.

[0116] An open reading frame (ORF) in this 2329 bp was retrieved, to find that the two kinds of protein were coded. Each amino acid sequence of these proteins was retrieved by the BLAST search, and as a result, an ORF (hereinafter referred to as bpmA) encoding a protein consisting of 395 amino acids having high homology to cytochrome P450 existed in the base 420 to base 1604 of the sequence No. 2. The bpmA had the highest homology (homology: 67.4%) to the amino acid sequence of the psmA isolated from the A-1544 strain and also had a relatively high homology (homology: 64.8%) to cytochrome P450 soy (Soy C) of *Streptomyces griseus*. It was considered from this fact that the bpmA highly possibly encoded hydroxylating enzyme of the cytochrome P-450 type.

[0117] Also, an ORF (hereinafter referred to as bpmB) encoding a protein having a high homology to ferredoxin of a 3Fe-4S type that existed just downstream (the base 1643 to base 1834 of the sequence No. 2) of the bpmA. The protein encoded by the bpmB consisted of 64 amino acids, and had the highest homology (81.0%) to the amino acid sequence of the psmB isolated from the A-1544 strain and a relatively higher homology (homology: 76.2%) to the amino acid sequence assumed to be that of ferredoxin just downstream of the amino acid sequence assumed to be cytochrome P450 (CYP105D5) of *Streptomyces coelicolor* A3(2). Therefore, it was considered that the bpmB served to transfer electrons and participated in hydroxylation together with the bpmA.

Example 8 Production of a transformant having the bpmA and the bpmB

(1) Preparation of a DNA fragment containing both the bpmA and the bpmB derived from the Mer-11107 strain

[0118] A primer 07-NdeF (sequence No. 14) obtained by adding a NdeI site to the 5' terminal and a primer 07-SpeR (sequence No. 15) obtained by adding a SpeI site to the 5' terminal were designed and produced on reference to the nucleotide sequence of the sequence No. 2 analyzed in Example 7: Next, these two types of primers (07-NdeF and 07-SpeR) and the Mer-11107 strain chromosome DNA obtained in Example 7(1) as a template, were used to run a PCR reaction. The PCR reaction was accomplished by repeating a two-stage reaction including denaturing carried out at 98°C for 20 seconds and annealing and elongation carried out at 68°C for 2 minutes 30 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and a PCR amplifier (T Gradient, Biometra Co.).

[0119] As a result, a DNA fragment (hereinafter referred to as a DNA fragment-D2) having a size of about 1.5 kbp and containing the psmA and the psmB was amplified. The DNA fragment-D2 was recovered from the PCR amplified reaction solution by SUPREC PCR (TAKARA HOLDINGS INC.).

(2) Architecture of a plasmid pTC-D07

[0120] pT7NS-CamAB (see WO03/087381) was digested by respective restriction enzymes NdeI and SpeI in a H

buffer solution (50 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 1 mM dithiothreitol and 100 mM NaCl) to obtain a plasmid digested products. Similarly, the DNA fragment-D2 obtained in the above (1) was digested by respective restriction enzymes NdeI and SpeI. The obtained digested product of the DNA fragment-D2 and the plasmid digested product were bound using DNA Ligation Kit ver. 2 (TAKARA HOLDINGS INC.). Thereby, a plasmid (referred to as a plasmid pTC-D07) about 9.5 kbp in size which was an architecture of a combination of the DNA fragment-D2 containing both the bpmA and the bpmB therein and the plasmid pT7NS-CamAB was formed.

(3) Preparation of E. coli transforming strain BL21 (DE3)/pTC-D07

[0121] Using the plasmid pTC-D07 prepared in the above (2), a competent cell (Novagen) of Colibacillus BL21 (DE3) was transformed. Thereby, E. coli BL21 (DE3)/pTC-D07 strain transformed by the plasmid pTC-D07 was obtained.

Example 9: Conversion of the macrolide compound 11107B into the 11107D by the E. coli transformant having the bpmA and the bpmB

[0122] The transformed E. coli BL21(DE3)/pTC-D07 strain obtained in Example 8(3) and a frozen seed of a BL21 (DE3)/pT7NS-CamAB strain were inoculated into a 15 mL test tube containing 3 mL of a L-broth medium (1.0% bactotripton, 0.5% yeast extract, 0.5% NaCl) containing 50 µg/mL of ampicillin and shake-cultured at 37°C for 20 hours. 500 µL of the seed culture broth was inoculated into a 250 mL Erlenmeyer flask containing 50 mL of a L-broth medium (1.0% bactotripton, 0.5% yeast extract, 0.5% NaCl) containing 50 µg/mL of ampicillin and shake-cultured at 32°C for 4 hours. Then, 50 µL of 100 mM IPTG (isopropyl-β-D-thiogalactopyranoside) and 50 µL of 80 mg/mL 5-aminolevulinic acid were successively added and the medium was shake-cultured at 32°C for 5 hours. The obtained culture broth was centrifuged (5000 rpm, 10 minutes) to collect the mycelia. The mycelia were then suspended in 1.75 mL of a 100 mM phosphate buffer solution (pH 6.1), to which were then added 250 µL of 80% glycerol and 12.5 µL of 40 mg/mL macrolide compound 11107B. The conversion reaction solution obtained in this manner was reacted at 28°C for 24 hours. 400 µL of the reaction solution was extracted with 600 µL of methanol and the extract was subjected to HPLC to measure each amount of macrolide compounds 11107B and 11107D. The results are shown in Table 6.

[0123] Also, the details of the condition of HPLC are shown below.

Analyzer: Shimadzu HPLC 10Avp

Column: Develosil ODS UG-3 (φ4.6 mm × 250 mm 3 µm)

Mobile phase: 45% to 55% methanol (0 to 5 minutes)

55% methanol (5 to 13 minutes)

55% to 70% methanol (13 to 17 minutes)

70% methanol (17 to 21 minutes)

45% methanol (21 to 25 minutes)

Flow rate: 1.2 mL/min.

Detection: UV 240 nm

Injection capacity: 5 µL

Column temperature: 40°C

Analyzing time: 25 minutes

Retention time: 11107B 12.2 minutes

11107D 4.2 minutes

Table 6

mg/L	BL21 (DE3)/pT7NS-CamAB	BL21 (DE3)/pTC-D07
11107B	162	156
11107D	0.00	0.78

[0124] As a result, the peak of the macrolide compound 11107D was not observed in the case of the E. coli BL21 (DE3)/pT7NS-CamAB strain used as a control, whereas the peak of the macrolide compound 11107D was obtained in the case of the BL21 (DE3)/pTC-D07 strain containing the psmA and psrB. This fact suggests that the bpmA and the bpmB participate in the conversion of the macrolide compound 11107B into the macrolide compound 11107D. Example 10 Determination of the nucleotide sequence of a gene derived from the A-1560 strain (FERM BP-10102)

(1) Preparation of a DNA of the A-1560 strain chromosome

[0125] The A-1560 strain was inoculated into a medium containing 1% of glucose, 0.4% of malt extract and 1% of yeast extract and cultured at 28°C for 3 days. The obtained culture broth was centrifuged at 3000 rpm for 10 minutes to collect the mycelia. A chromosome DNA was prepared using Blood & Cell Culture kit (QIAGEN Co.) from the mycelia.

(2) Cloning of a partial sequence of a DNA encoding a protein having the activity in hydroxylating the 16-position of the macrolide compound 11107

[0126] Mix primers (5Dm-3F (sequence no. 4) and 5Dm-2R (sequence No. 16) were designed and produced on reference to the amino acid sequence estimated as that of the cytochrome P450 (CYP105D5) of *Streptomyces coelicolor* A3(2).

[0127] In order to promote reactivity taking the fluctuation of a codon into account, mixed bases S (=C+G) and Y (=C+T) were used.

[0128] Next, these two types of primers (5Dm-3F and 5Dm-2R) and the A-1560 strain chromosome DNA obtained in the above (1) as a template, were used to run a PCR reaction. The PCR reaction was accomplished by repeating a three-stage reaction including denaturing carried out at 98°C for 20 seconds, annealing carried out at 50°C for 2 minutes and elongation carried out at 68°C for 30 seconds 35 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and a PCR amplifier (T Gradient, Biometra Co.). As a result, a DNA fragment (hereinafter referred to as a DNA fragment-A3) having a size of about 750 bp was amplified. It is highly possible that this DNA fragment-A3 is a part of the DNA encoding a protein having hydroxylating activity. The DNA fragment-A3 amplified by a PCR reaction was recovered from the reaction solution by SUPREC PCR (TAKARA HOLDINGS INC.).

[0129] In order to obtain the DNA fragment-A3 in an amount enough to analyze the nucleotide sequence of the obtained DNA fragment-A3, the DNA fragment-A3 was bound with a plasmid vector pT7Blue T (Novagen Co.) by using DNA Ligation kit ver.2 (TAKARA HOLDINGS INC.) to transform *E. coli* JM109 strain (Stratagene Co.). Thereafter, the transformed *E. coli* was selected using a L-broth agar media (1.0% bactotripton, 0.5% yeast extract, 0.5% NaCl, 1.5% agar) containing ampicillin (50 µg/mL), X-gal (5-bromo-4-chloro-3-indolyl-β-D-galactoside; 40 µg/mL) and IPTG (isopropyl-β-D-thiogalactopyranoside; 100 µM). The colony of the transformed *E. coli* thus isolated was cultured in a L-broth liquid medium (1% bactotripton, 0.5% yeast extract, 0.5% NaCl) containing ampicillin (50 µg/mL). A plasmid DNA was separated from the mycelia of the proliferated transformed *E. coli* and purified by using a plasmid purifying kit (QIA filter Plasmid Midi Kit, QIAGEN Co.), to obtain enough amount of the DNA fragment-A3.

(3) Analysis of the nucleotide sequence of the cloned DNA fragment-A3

[0130] The nucleotide sequence of the DNA fragment-A3 obtained in the above (2) was analyzed using a DNA nucleotide sequence analyzer (PE Biosystems 377XL) according to a dye terminator cycle sequence method. As the result of the nucleotide sequence analysis, it was clarified that the DNA fragment-A3 amplified by a PCR reaction had an exact size of 741 bp though it had been found to have a size of about 750 bp by the above measurement using electrophoresis (see the nucleotide sequence 616 to nucleotide sequence 1356 of the sequence No. 3). Since DNA sequences corresponding to the two types of primers used in the above PCR reaction were found at both ends of the above cloned 741 bp DNA sequence, it was clarified that the DNA fragment-A3 was singularly amplified by these two types of primers (5Dm-3F and 5Dm-2R) in the above PCR reaction.

(4) Analysis of the neighboring region of the DNA fragment-A3

[0131] As mentioned above, the partial sequence of the DNA encoding a protein having hydroxylating activity derived from the A-1560 strain was determined. Therefore, the amplification, cloning and sequence analysis of the nucleotide sequence in the neighboring region extending from the upstream side to downstream side of the cloned fragment were accomplished by an inverse PCR method (Cell Technology vol. 14, p.591-593, 1995).

Specifically, the A-1560 strain chromosome DNA (see the above (1)) was digested by a restriction enzyme BamHI in a K buffer solution (50 mM Tris-HCl, pH 8.5, 10 mM MgCl₂, 1 mM dithiothreitol and 100 mM KCl), by a restriction enzyme KpnI in a L buffer solution (10 mM Tris-HCl, pH 7.5, 10 mM MgCl₂ and 1 mM dithiothreitol) and by a restriction enzyme Sall in a H buffer solution (50 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 1 mM dithiothreitol and 100 mM NaCl) respectively. The obtained each DNA fragment cut by the restriction enzymes was self-circularized using DNA Ligation Kit ver.2 (TAKARA HOLDINGS INC.).

[0132] On the other hand, primers (5PIN-2F (sequence No. 17) and 6PIN-2R (sequence No. 7)) were designed and produced based on the nucleotide sequence of the DNA fragment-A3.

[0133] Next, these two primers (5PIN-2F and 6PIN-2R) and the above self-circularized A-1560 strain chromosome

DNA as a template, were used to run a PCR reaction. In the PCR reaction, the cycle of a two-stage reaction involving denaturing carried out at 98°C for 20 seconds and annealing and elongation carried out at 68°C for 5 minutes was repeated 35 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and PCR amplifier (T Gradient, Biometra Co.).

[0134] As a result, a DNA fragment (DNA fragment-B3) about 4.5 kbp in size, a DNA fragment (DNA fragment-C3) about 3.0 kbp in size and a DNA fragment (DNA fragment-D3) about 1.7 kbp in size were amplified. It was highly possible that these DNA fragments were a DNA encoding a protein having hydroxylating activity and a DNA having a DNA sequence including those in the upstream and downstream regions of the former DNA.

[0135] The DNA fragment-B3, the DNA fragment-C3 and the DNA fragment-D3 were recovered from the PCR amplified reaction solution by SUPREC PCR (TAKARA HOLDINGS INC.). Next, as to the obtained DNA fragment-B3, DNA fragment-C3 and DNA fragment-D3, in order to obtain each DNA fragment in an amount enough to analyze the nucleotide sequence of the obtained DNA fragment, a plasmid vector pT7Blue T (Novagen Co.), DNA Ligation kit ver.2 (TAKARA HOLDINGS INC.), E. coli JM109 strain and a plasmid purifying kit (QIA filter Plasmid Midi Kit, QIAGEN Co.) were used in the same manner as the above (2), to obtain enough amount of each DNA fragment.

(5) Analysis of each nucleotide sequence of the DNA fragment-B3 (about 4.5 kbp in size), the DNA fragment-C3 (about 3.0 kbp in size) and the DNA fragment-D3 (about 1.7 kbp in size)

[0136] Each nucleotide sequence of the DNA fragment-B3, DNA fragment-C3 and DNA fragment-D3 obtained in the above (4) was analyzed using a DNA nucleotide sequence analyzer (PE Biosystems 377XL) according to a dye terminator cycle sequence method. The nucleotide sequence was thus analyzed to obtain the information of the nucleotide sequence of 1860 bp shown in the sequence No. 3 from each sequence of the DNA fragment-B3, DNA fragment-C3 and DNA fragment-D3.

[0137] An open reading frame (ORF) in this 1860 bp was retrieved, to find that the two kinds of protein were encoded. Each amino acid sequence of these proteins was retrieved by the BLAST search, and as a result, an ORF (hereinafter referred to as tpmA) encoding a protein consisting of 404 amino acids having high homology to cytochrome P450 existed in the base 172 to base 1383 of the sequence No. 3. The tpmA had the highest homology (homology: 77.4%) to the amino acid sequence assumed to be that of cytochrome P450 (CYP105D5) of *Streptomyces coelicolor* A3 (2) and also a high homology (homology: 76.6%) to the amino acid sequence of the psmA isolated from the A-1544 strain. It was considered from this fact that the tpmA was highly possibly a gene encoding hydroxylating enzyme of the cytochrome P-450 type.

[0138] Also, an ORF (hereinafter referred to as tpmB) encoding a protein having a high homology to ferredoxin of a 3Fe-4S type existed just downstream (the base 1399 to base 1593 of the sequence No. 3) of the tpmA. The protein encoded by the tpmB consisted of 65 amino acids, and had the highest homology (81.0%) to the amino acid sequence of the psmB isolated from the A-1544 strain and also a high homology (homology: 82.5%) to the amino acid sequence assumed to be that of ferredoxin just downstream of the amino acid sequence assumed to be cytochrome P450 (CYP105D5) of *Streptomyces coelicolor* A3 (2). Therefore, it was considered that the tpmB served to transfer electrons and coded ferredoxin participating in hydroxylation together with the tpmA.

Example 11 Production of a transformant having the tpmA and the tpmB

(1) Preparation of a DNA fragment containing both the tpmA and the tpmB derived from the A-1560 strain

[0139] A primer tpm-NdeF (sequence No. 18) obtained by adding a NdeI site to the 5' terminal and a primer tpm-SpeR (sequence No. 19) obtained by adding a SpeI site to the 5' terminal were designed and produced on reference to the nucleotide sequence of the sequence No. 3 analyzed in Example 10. Next, these two types of primers (tpm-NdeF and tpm-SpeR) and the A-1560 strain chromosome DNA obtained in Example 10(1) as a template, were used to run a PCR reaction. The PCR reaction was accomplished by repeating a two-stage reaction including denaturing carried at 98°C for 20 seconds and annealing and elongation carried out at 68°C for 2 minutes 30 times by using Takara LA Taq (TAKARA HOLDINGS INC.) and a PCR amplifier (T Gradient, Biometra Co.).

[0140] As a result, a DNA fragment (hereinafter referred to as a DNA fragment-E3) having a size of about 1.5 kbp and containing the tpmA and the tpmB was amplified. The DNA fragment-E3 was recovered from this PCR amplified reaction solution by SUPREC PCR (TAKARA HOLDINGS INC.).

(2) Architecture of a plasmid pTC-tpmAB

[0141] pT7NS-CamAB (see WO03/087381) was digested by respective restriction enzymes NdeI and SpeI in a H buffer solution (50 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 1 mM dithiothreitol and 100 mM NaCl) to obtain plasmid digested products. Similarly, the DNA fragment-E3 obtained in the above (1) was digested by respective restriction enzymes NdeI

and Spel. The obtained digested product of the DNA fragment-E3 and the plasmid digested product were bound using DNA Ligation Kit ver. 2 (TAKARA HOLDINGS INC.). Thereby, a plasmid (referred to as a plasmid pTC-tpmAB) about 9.5 kbp in size which was an architecture of a combination of the DNA fragment-E3 containing both the tpmA and the tpmB therein and the plasmid pT7NS-CamAB was formed.

(3) Preparation of E. coli transforming strain BL21 (DE3)/pTC-tpmAB

[0142] Using the plasmid pTC-tpmAB prepared in Example 11(2), a competent cell (Novagen) of Colibacillus BL21 (DE3) was transformed, to give E. coli BL21 (DE3)/pTC-tpmAB strain transformed by the plasmid pTC-tpmAB.

Example 12: Conversion of the 11107B into the 11107D by the E. coli transformant having the tpmA and the tpmB

[0143] The transformed E. coli BL21(DE3)/pTC-tpmAB strain obtained in the above (3) and a frozen seed of a BL21 (DE3) /pT7NS-CamAB strain were inoculated into a 15 mL test tube containing 3 mL of a L-broth medium (1.0% bactotripton, 0.5% yeast extract, 0.5% NaCl) containing 50 µg/mL of ampicillin and shake-cultured at 37°C for 20 hours. 500 µL of the seed culture broth was inoculated into a 250 mL Erlenmeyer flask containing 50 mL of a L-broth medium (1.0% bactotripton, 0.5% yeast extract, 0.5% NaCl) containing 50 µg/mL of ampicillin and shake-cultured at 32°C for 4 hours. Then, 50 µL of 100 mM IPTG (isopropyl-β-D-thiogalactopyranoside) and 50 µL of 80 mg/mL 5-aminolevulinic acid were successively added thereto, and the medium was shake-cultured at 32°C for 5 hours. The obtained culture broth was centrifuged (5000 rpm, 10 minutes) to collect the mycelia. The mycelia were suspended in 1.75 mL of a 100 mM phosphate buffer solution (pH 6.1), and 250 µL of 80% glycerol and 12.5 µL of 40 mg/mL macrolide compound 11107B were added thereto. The conversion reaction solution obtained in this manner was reacted at 28°C for 24 hours. 400 µL of the reaction solution was extracted with 600 µL of methanol and the extract was subjected to HPLC to measure each amount of macrolide compounds 11107B and 11107D. The results are shown in Table 7. Also, the details of the condition of HPLC are shown below. Analyzer: Shimadzu HPLC 10Avp
Column: Develosil ODS UG-3 (φ4.6 mm × 250 mm 3 µm)
Mobile phase: 45% to 55% methanol (0 to 5 minutes)
55% methanol (5 to 13 minutes)
55% to 70% methanol (13 to 17 minutes)
70% methanol (17 to 21 minutes)
45% methanol (21 to 25 minutes)
Flow rate: 1.2 mL/min.
Detection: UV 240 nm
Injection capacity: 5 µL
Column temperature: 40°C
Analyzing time: 25 minutes
Retention time: 11107B 12.2 minutes
11107D 4.2 minutes


Table 7

mg/L	BL21(DE3)/pT7NS-CamAB	BL21(DE3)/pTC-tpmAB
11107B	141	128
11107D	0	18

[0144] As a result, the peak of the macrolide compound 11107D was not observed in the case of the E. coli BL21 (DE3)/pT7NS-CamAB strain used as a control, whereas the peak of 11107D was obtained in the case of the BL21(DE3) /pTC-tpmAB strain containing the tpmA and tpmB. This fact suggests that the tpmA and the tpmB participate in the conversion of 11107B into 11107D.

Example 13: Conversion of 11107H into 11107CB represented by the following formulae by a self-cloning strain

[0145]



(1) Preparation of a transformant reaction solution

[0146] A medium containing 2.0% of Stabilose, 2.0% of glucose, 2.0% of a soybean meal (Honen Soypro), 0.5% of yeast extract and 0.32% of CaCO_3 and having a pH of 7.4 was prepared. A 250 mL Erlenmeyer flask was charged with 25 mL of the medium, which was then sterilized under heating at 121°C for 20 minutes and thiostrepton was added to the medium such that its final concentration was 25 mg/L. Then, 1% of an A-1544/pJDMG strain from frozen seed was inoculated to culture the seed at 28°C and 220 rpm for 3 days. 1% of the seed culture broth was added in a medium having the same composition to carry out main culturing at 28°C and 220 rpm for 2 days. After the main culturing was finished, mycelia were collected from the culture broth by centrifugation and suspended in 20 mL of phosphate buffer solution having a pH of 6.5. The substrate 11107H (100 g/L DMSO solution) was added in this mycelia suspended solution such that its final concentration was 2000 mg/L to run a conversion reaction at 28°C and 220 rpm for 16 hours.

(2) Isolation of amacrolide compound 11107CB from a transformant reaction solution


[0147] Mycelia were isolated from a conversion reaction solution (in an amount corresponding to 6 flasks) obtained from the same operation, by centrifugation and the centrifuged supernatant was extracted with the equal amount of ethyl acetate twice. The extract was concentrated and then the residue was purified by thin layer chromatography (MERCK Silicagel 60 F254' 0.5 mm, developing solution: toluene:acetone=1:1), to obtain 119.5 mg of 11107CB.

¹H-NMR spectrum (CD_3OD , 500MHz): δ ppm (integral, multiplicity, coupling constant J(Hz)):

0.81(3H,d,J=6.7Hz), 0.89(3H,d,J=7.0), 0.94(3H,t,J=7.4Hz), 1.25(3H,s), 1.30-1.20(1H,m), 1.33(3H,s), 1.55-1.40(2H,m), 1.65(1H,dd,J=6.3,14.0Hz), 1.75(3H,s), 1.88(1H,dd,J=5.4,14.0Hz), 2.07(3H,s), 2.68-2.40(4H,m), 2.89(1H,m), 3.51(1H,m), 4.51(1H,m), 4.97(1H,d,J=8.6Hz), 4.99(1H,d,J=9.3Hz), 5.30(1H,dd,J=9.7,15.2Hz), 5.52(1H,dd,J=9.4,15.2Hz), 5.58(1H,dd,J=1.9,15.5Hz), 5.78(1H,dd,J=2.8,15.5Hz), 5.85(1H,d,J=15.3Hz), 6.07(1H,d,J=11.0Hz), 6.51(1H,dd,J=11.0,15.3Hz)

Example 14: Conversion of 11107L into 11107CG represented by the following formulae respectively by a self-cloning strain

[0148]



(1) Preparation of a transformant reaction solution

[0149] A medium containing 2.0% of stabilose, 2.0% of glucose, 2.0% of a soybean meal (Honen Soypro), 0.5% of yeast extract and 0.32% of CaCO_3 and having a pH of 7.4 was prepared. A 250 mL Erlenmeyer flask was charged with 25 mL of the medium, which was then sterilized under heating at 121°C for 20 minutes and thiostrepton was added to the medium such that its final concentration was 25 mg/L. Then, 1% of an A-1544/pIJDMG strain from frozen stock was inoculated to cultivate the seed culture at 28°C and 220 rpm for 3 days. 1% of this seed culture broth was added in a medium having the same composition to carry out main cultivation at 28°C and 220 rpm for 2 days. After the main cultivation was finished, mycelia were collected from the culture broth by centrifugation and suspended in 20 mL of phosphate buffer solution having a pH of 6.5. The substrate 11107L (100 g/L DMSO solution) was added to this mycelia suspension solution such that its final concentration was 1600 mg/L to run a conversion reaction at 28°C and 220 rpm for 16 hours.

(2) Isolation of a macrolide compound 11107CG from a transformant reaction solution

[0150] Mycelia were isolated from the conversion reaction solution by centrifugation and the centrifuged supernatant was extracted with the equivalent amount of ethyl acetate twice. The extract layers were concentrated and then the residue was purified by thin layer chromatography (MERCK Silicagel 60 F254, 0.25 mm, developing solution: toluene: acetone=1:1), to obtain 25 mg of 11107CG.

ESI-MS m/z 633 ($M+\text{Na}$)⁺

¹H-NMR spectrum (CD_3OD , 500MHz) : δ ppm (integral, multiplicity, coupling constant J (Hz)):

0.88(3H,d,J=6.7Hz), 0.90(3H,d,J=7.0Hz), 0.94(3H,d,J=7.4Hz), 1.18(3H,s), 1.30-1.20(1H,m), 1.39,(3H,s), 1.56-1.40(2H,m), 1.66(1H,dd,J=6.2,14.0Hz), 1.79-1.69(2H,m), 1.81(3H,d,J=1.0Hz), 1.86(1H,dd,J=5.4,14.0Hz), 2.05(3H,s), 2.08(3H,s), 2.52(1H,dd,J=4.2,15.2Hz), 2.64-2.55(1H,m), 2.67(1H,dd,J=2.2,7.9Hz), 2.78(1H,dd,J=3.0,15.2Hz), 2.90(1H,dt,J=2.2,5.6Hz), 3.52(1H,dt,J=4.4,8.8Hz), 3.75(1H,m), 4.98(1H,dd,J=2.8,11.3Hz), 5.08(1H,d,J=9.7Hz), 5.13(1H,d,J=9.6Hz), 5.61(1H,dd,J=9.9,15.2Hz), 5.75(1H,dd,J=9.7,15.2Hz), 5.88(1H,d,J=15.3Hz), 6.13(1H,d,J=11.0Hz), 6.54(1H,dd,J=11.0,15.3Hz)

Industrial applicability

[0151] A 12-membered macrolide compound which has hydroxyl group at the 16-position and is excellent in antitumor

activity and stability in an aqueous solution can be produced efficiently by using a transformant obtained by transformation using a plasmid carrying the DNA of the present invention.

5

10

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

5 <110> Mercian Corporation
 <110> Eisai Co., Ltd
 <120> DNA related to hydroxylation of macrolide compounds
 10 <130> 04063PCT
 <150> JP 2003-396828
 <151> 2003-11-27
 <160> 19
 15 <210> 1
 <211> 3793
 <212> DNA
 <213> Streptomyces sp.
 20 <220>
 <221> CDS
 <222> (1322).. (2548)
 <220>
 25 <221> CDS
 <222> (2564).. (2761)
 <400> 1
 ctgcagctcg acgtcgsggt cggacttcac gtigaagtac cagaccggat gcttggcgcc 60
 accgcccagc gaggcaccg ccgcgtaact cccctcgtgc tcgaccgcga tcagcgsgct 120
 cttagcgatc tticccctgc gcgcgccccg ggtggtagc acgatgacce gcagcccgg 180
 gtcccgcagc gtggcgccct tggcgcccc ggaactctcg tacagctcga cctgctcgcg 240
 caccctctgc gtcggcgctg gctcgtacgc gccctcaagt ggcaagggat ccgtctcctt 300
 cgtcggtccg gcggaiggtg ctccggacgg tcccaactcc cgcggccgcc cggatcatcc 360
 gtaccgcatg ccccttcgcc cgagcgggtg atcacccttc cggccatccg gtcgtccgca 420
 ccgcgagcac caggatcacg gcgctggaga gcaggcgctt gaccagccgc ccccggtggc 480
 ccctcagggc gcgacccagc agcgcgcccc cgcgcgcccag cagtagctgc cagctcgcgg 540
 acgcggcgaa ggccgcccgc gcgaacacgc cccgtttcag cggccgtgcc gcaccggcgg 600
 cgccgtctgc gaggaccagc gccacgaagt agaccaccgt catgggattg agcagggtga 660
 tcccagaaag gccgagataa gcccttgcgc gccttgaac cggccgttcc gggcgsgtgg 720
 tgagccgatg ggcgcggtac tgcgcagggc cgagcagcgc cgcgcgagc gcgagcaccg 780
 cgaggaccag cgcgagggcc cagcgcagcg gctccagcac cgcgcgagc tgcgcgca 840
 ggccggcgcc gccacggctc gcgagcagcg cgtacagccc gtcggccgtg gcgacgcca 900
 gcgcgcgcca ggcgcgsgtg cgcagcgagg tgcggcgctt gaggagagacc agataggtcc 960
 cgaccgcgcc gacgggcacc gcgatgccgt acccgcgag caggcccgcg agcagcgcg 1020
 ccgtcacggc cgtgcgggac tggctcctcc gggacggcg gggctgctgt cggcccgca 1080
 ccgcggggcg ggtggcagcg ggcgicggca ggaggagggc tctaggaggc atggccgat 1140
 cctggggcg ccgcgcccgc accggcaaat gaattacggc gcgttcagc ccccgcccg 1200
 ctgcctcttc ggccactica ccgcgtacgg ccatctgcc gaacttgcg tcgccccata 1260
 ggtgcctcgg gcatctaatg aagatcgcca cgacgcacct ctctgtcgc gaggctttc 1320
 c atg acg gaa ctg acg gac atc acc gcc cgg acc cgg gcc gaa 1366
 Met Thr Glu Leu Thr Asp Ile Thr Gly Pro Gly Thr Pro Ala Glu
 1 5 10 15
 ccc gtc gca ttc ccc cag gac cgc acc tgc ccc tac cac ccc ccc acc 1414
 Pro Val Ala Phe Pro Gln Asp Arg Thr Cys Pro Tyr His Pro Pro Thr
 20 25 30
 gga tac ggc ccg ctg cgc gac ggg cgc agc ctg tcc cgc gtc acc ctc 1462
 Gly Tyr Gly Pro Leu Arg Asp Gly Arg Ser Leu Ser Arg Val Thr Leu
 35 40 45
 ttc gac ggc cgc gag gtc tgg atg gtc acc ggc cac gcc acc gcc cgc 1510
 Phe Asp Gly Arg Glu Val Trp Met Val Thr Gly His Ala Thr Ala Arg

EP 1 705 247 A1

	50	55	60	
5	gcg ctg ctc gcg gac ccc cgg ctg tcc acc gac cgc acc ctc ccg ggc Ala Leu Leu Ala Asp Pro Arg Leu Ser Thr Asp Arg Thr Leu Pro Gly	1558		
	65	70	75	
	ttc ccc gtg ccc acg gcc cgc ttc gcg gcc gtc cgc gac cgg cgg gtg Phe Pro Val Pro Thr Ala Arg Phe Ala Ala Val Arg Asp Arg Arg Val	1606		
	80	85	90	95
10	gcg ctg ctc ggc gtg gac cgc cgc gtc cac cag acc cag cgg cgg atg Ala Leu Leu Gly Val Asp Asp Pro Val His Gln Thr Gln Arg Arg Met	1654		
	100	105	110	
	atg atc ccc tgc ttc acc ctc aag cgc ggc gcc ggg ctg cgg ccc acc Met Ile Pro Ser Phe Thr Leu Lys Arg Ala Ala Gly Leu Arg Pro Thr	1702		
	115	120	125	
15	atc cag cgg acc gtc gac ggg ctg ctg gac gcg atg atc gag aag ggg Ile Gln Arg Thr Val Asp Gly Leu Leu Asp Ala Met Ile Glu Lys Gly	1750		
	130	135	140	
	ccg cgg gcc gag ctg gtc tcc gcc ttc gcc ctg ccc gtc ccc tgg gtc Pro Pro Ala Glu Leu Val Ser Ala Phe Ala Leu Pro Val Pro Ser Val	1798		
	145	150	155	
20	gtc atc tgc ggc ctg ctc ggc gtc cgg tac gcc gac cac gag ttc ttc Val Ile Cys Gly Leu Leu Gly Val Pro Tyr Ala Asp His Glu Phe Phe	1846		
	160	165	170	175
	gag gaa cag tcc cgc acg ctg ctg cgc ggt ccc acg gcc gcc gac tgc Glu Glu Gln Ser Arg Thr Leu Leu Arg Gly Pro Thr Ala Ala Asp Ser	1894		
	180	185	190	
25	caa ggg gcg cgc gag cgg ctg gag gag tac ctg ggc ggg ctg atc gac Gln Gly Ala Arg Glu Arg Leu Glu Glu Tyr Leu Gly Gly Leu Ile Asp	1942		
	195	200	205	
	gac aag gag cgg cag gcc gaa ccc ggc gac ggc gtc ctg gac gac ctg Asp Lys Glu Arg Gln Ala Glu Pro Gly Asp Gly Val Leu Asp Asp Leu	1990		
	210	215	220	
30	gtc cac cag cgg ctg cgc acc ggc gag ctg gac cgg cgc gac gtc gtc Val His Gln Arg Leu Arg Thr Gly Glu Leu Asp Arg Arg Asp Val Val	2038		
	225	230	235	
	gcg ctg gcc gtc atc ctg ctc gtc gcc ggg cac gag acg acc gcc aac Ala Leu Ala Val Ile Leu Leu Val Ala Gly His Glu Thr Thr Ala Asn	2086		
	240	245	250	255
35	atg atc tcc ctc ggc acc tac acg ctg ctg cgg cac ccc ggc cgg ctg Met Ile Ser Leu Gly Thr Tyr Thr Leu Leu Arg His Pro Gly Arg Leu	2134		
	260	265	270	
	gcc gag ctg cgc gcc gac cgc ggc ctg ctg ccc gcc gcc gtc gag gag Ala Glu Leu Arg Ala Asp Pro Ala Leu Leu Pro Ala Ala Val Glu Glu	2182		
	275	280	285	
40	ctg atg cgg atg ctc tgc atc gcg gac ggg ctg ctg cgc ctg gcc ctg Leu Met Arg Met Leu Ser Ile Ala Asp Gly Leu Leu Arg Leu Ala Leu	2230		
	290	295	300	
	gag gac atc gag atc gcc ggc gcc acg atc cgg gcc ggc gag ggc gtc Glu Asp Ile Glu Ile Ala Gly Ala Thr Ile Arg Ala Gly Glu Gly Val	2278		
	305	310	315	
45	ctg ttc tcc acc tgc ctg atc aac cgc gac gag tcc gtc ttc gac gac Leu Phe Ser Thr Ser Leu Ile Asn Arg Asp Glu Ser Val Phe Asp Asp	2326		
	320	325	330	335
	ccc gac acc ctg gac ttc cac cgc tcc acc cgc cac cac gtc gcc ttc Pro Asp Thr Leu Asp Phe His Arg Ser Thr Arg His His Val Ala Phe	2374		
	340	345	350	
50	ggt ttc ggc atc cac cag tgc ctg ggc cag aac ctg gcc cgc gcc gag Gly Phe Gly Ile His Gln Cys Leu Gly Gln Asn Leu Ala Arg Ala Glu	2422		
	355	360	365	
	ctg gag atc gcc ctg ggc acg ctc ctg gag cgg ctc ccc ggc ctc cgg Leu Glu Ile Ala Leu Gly Thr Leu Leu Glu Arg Leu Pro Gly Leu Arg	2470		
55	370	375	380	

5	<p> ctg gcc gcg ccc gcc gag gag atc ccg ttc aaa ccc gcc gac acg atc Leu Ala Ala Pro Ala Glu Glu Ile Pro Phe Lys Pro Gly Asp Thr Ile 385 390 395 </p> <p> cag ggg atg ctg gaa ctc ccc gtg acc tgg taa gaagctctgg tc atg cac Gln Gly Met Leu Glu Leu Pro Val Thr Trp Met His 400 405 410 </p> <p> atc gac atc gac aag gac cgc tgc atc ggc gcc ggc cag tgc gcg ctg Ile Asp Ile Asp Lys Asp Arg Cys Ile Gly Ala Gly Gln Cys Ala Leu 415 420 425 </p> <p> gcc gcc ccg gcc gtc ttc acc cag gac gac gac ggc tac agc acc ctg Ala Ala Pro Gly Val Phe Thr Gln Asp Asp Asp Gly Tyr Ser Thr Leu 430 435 440 </p> <p> ctc ccc gcc gcg gag gac gcc ggg gcc gac ccg atg gtc ccg gag gcc Leu Pro Gly Arg Glu Asp Gly Gly Gly Asp Pro Met Val Arg Glu Ala 445 450 455 </p> <p> gcc cgc gcc tgc ccg gtc agc gcc atc ccg gtc acc gaa ccg gcc gcc Ala Arg Ala Cys Pro Val Ser Ala Ile Arg Val Thr Glu Pro Ala Gly 460 465 470 475 </p>	<p>2518</p> <p>2569</p> <p>2617</p> <p>2665</p> <p>2713</p> <p>2761</p>
20	<p>tga ggccggggccc ggccggccgc gccgcctgcc gggaccgcgc ttccagttc agtagg</p>	<p>2820</p>
25	<p> gtcgtgcgat gacctcacag gccgggaagc ctttctcta cgtcgtcgtc tgcgcggccg ggaccgccgc cggagtcacc acgcgatcgc gcgccgccca gccgcggggc tggaggatgg gggtcttggc caccgcggtg gcgatgggcg ggttcttcga caccgctgcg gtccaggaga tgaccggccg gcccatccgc tggccttggc gctcgccgcg cgatccgcgc ccgttccgc cgccggccgc cgtggttggt gcgccgccca cttcaaacac cgtcaacaag tggcggccg gtctcgccga cagcctgcgc gtccgcacgc tctcgaggc gccggccctc ggcgtgccga tcccttccct gccctgcgt gcgacgcgc tggccgccca ccccgctac cggaggggcc ttctccgct gcctgggatg ggcgtccgt tggcgagcc gtacgcggcg ccgccgggg aggacggcga ggcggacgcg gcacggcccg gttcgcctg ggagaacgcc ctggacctgc tggagcggcg ctgaacctcg tccccgacc gtaggccctg tctgacactg tcagacaggc cttaacggca ggtcagccg gcccggccca gcatccgcc ggtgtagagg tcttggcccc gcggcagcca gttagccagc cggagacca ccglggagca gtcaggcccg acggtgacgc ggaccttac cgtctcgga cggccggct gcagcgcgt cagcgcgcag tccaggaggt acggagccg ggtcttcgag gtaccggccg accagcgggt tgacgcagcg ggcgtcgtcc gtggcgaicc gcacccgggt gcccgcccg ccgatgagtc cagccgggc gctgcgctcg ccctcgtgc tgtcccggcg gacctgtag gtccgctgg tegtgcgtc gccccgcagg gtgtccgctc gac </p>	<p>2880</p> <p>2940</p> <p>3000</p> <p>3060</p> <p>3120</p> <p>3180</p> <p>3240</p> <p>3300</p> <p>3360</p> <p>3420</p> <p>3480</p> <p>3540</p> <p>3600</p> <p>3660</p> <p>3720</p> <p>3780</p> <p>3793</p>
30		
35		

40 <210> 2
 <211> 2329
 <212> DNA
 <213> Streptomyces sp.

45
 <220>
 <221> CDS
 <222> (420).. (1604)

<220>
<221> CDS
<222> (1643).. (1834)

[illegible]

	cgtagccatc tcacacacga gcaactcag ccacttgaga ctctacagg aggaatc	419
5	gtg acc gaa gcc atc ccc tac ttt cag aac cgc acc tgt ccc tac cac Val Thr Glu Ala Ile Pro Tyr Phe Gln Asn Arg Thr Cys Pro Tyr His	467
	1 5 10 15	
	ccg ccc gcc gcc tat cag cca ctg cgc ggg gcc ggc ccg ctg agc cat Pro Pro Ala Ala Tyr Gln Pro Leu Arg Gly Ala Gly Pro Leu Ser His	515
10	20 25 30	
	gtc acc ttc tac gac ggc cgg aag gtg tgg ggc gtc acc ggc cac ccc Val Thr Phe Tyr Asp Gly Arg Lys Val Trp Ala Val Thr Gly His Pro	563
	35 40 45	
	gag gca cgg ggc ctg ctg acc gac cag cga ctc tcc gcc gac cgg cag Glu Ala Arg Ala Leu Leu Thr Asp Gln Arg Leu Ser Ala Asp Arg Gln	611
15	50 55 60	
	aac ccg gcc ttc ccg gtc ccc ttc gaa cgc ttc ggc gcc atc cgc cgg Asn Pro Ala Phe Pro Val Pro Phe Glu Arg Phe Ala Ala Ile Arg Arg	659
	65 70 75 80	
	gtc cgg acc ccg ctg atc ggg gtc gac gac ccg gag cac aac acc cag Val Arg Thr Pro Leu Ile Gly Val Asp Asp Pro Glu His Asn Thr Gln	707
20	85 90 95	
	cgc cgg atg ctg atc ccc agc ttc agc ctc aag cgg acc gcc gca ctg Arg Arg Met Leu Ile Pro Ser Phe Ser Leu Lys Arg Thr Ala Ala Leu	755
	100 105 110	
	cgg ccg gag atc cag cgg atc gtc gac ggg ctg ctg gac cgg atg ctg Arg Pro Glu Ile Gln Arg Ile Val Asp Gly Leu Leu Asp Arg Met Leu	803
25	115 120 125	
	gat cag ggc ccg ccc acc gag ctg gtc tcc ggc ttc gcc ctg ccg gtc Asp Gln Gly Pro Pro Thr Glu Leu Val Ser Ala Phe Ala Leu Pro Val	851
	130 135 140	
	ccg tgg atg gtc atc tgc gca ctg ctg gga gtc tca tac gcc gac cat Pro Ser Met Val Ile Cys Ala Leu Leu Gly Val Ser Tyr Ala Asp His	899
30	145 150 155 160	
	gag ttc ttc gag gag gag tcc cgc cgc atc ctg cgc gcc cgg tgg gcc Glu Phe Phe Glu Glu Glu Ser Arg Arg Ile Leu Arg Gly Arg Ser Ala	947
	165 170 175	
	gag gag ggc gag gac gcc cgg ctg aag ctg gag gag tac ttc acc ggg Glu Glu Ala Glu Asp Ala Arg Leu Lys Leu Glu Glu Tyr Phe Thr Gly	995
35	180 185 190	
	ctg atc gcc ggc aag gag aag aac cgc ggc gac ggg ctg ctg gac gag Leu Ile Ala Ala Lys Glu Lys Asn Pro Gly Asp Gly Leu Leu Asp Glu	1043
	195 200 205	
	ctg atc gag gac cgg ctg cgg acc ggc ggc ctc acc cgc gac gag ctg Leu Ile Glu Asp Arg Leu Arg Thr Gly Ala Leu Thr Arg Asp Glu Leu	1091
40	210 215 220	
	gtc cgg ctc gcc atg atc ctg ctg gtc gcc ggc cat gag acc acc gcc Val Arg Leu Ala Met Ile Leu Leu Val Ala Gly His Glu Thr Thr Ala	1139
	225 230 235 240	
	aac atg atc tgg ctg ggc acc ttc acc ctg ctg gac cac ccc gag cag Asn Met Ile Ser Leu Gly Thr Phe Thr Leu Leu Asp His Pro Glu Gln	1187
45	245 250 255	
	ctg ggc cag ctc aag gcc gac gag ggc ctg atg ccg gcc gcc atc gag Leu Ala Gln Leu Lys Ala Asp Glu Gly Leu Met Pro Ala Ala Ile Glu	1235
	260 265 270	
	gag ctg ctg cga ttc ctg tcc atc cgc gac ggc ctg ctg cgg gtc ggc Glu Leu Leu Arg Phe Leu Ser Ile Ala Asp Gly Leu Leu Arg Val Ala	1283
50	275 280 285	
	acg gag gac atc gag atc ggc ggt cag gtc atc cgg gcc gac gac ggc Thr Glu Asp Ile Glu Ile Gly Gly Gln Val Ile Arg Ala Asp Asp Ala	1331
	290 295 300	
55	gtc ctg ttc ccc gcc tca ctg atc aac cgg gac gag gcc gcc tat ccg Val Leu Phe Pro Ala Ser Leu Ile Asn Arg Asp Glu Ala Ala Tyr Pro	1379

305 310 315 320
 gca ccc gac gag ctg gac ctc ggc cgt tgc gcc cgc cat cac gtg gcg 1427
 Ala Pro Asp Glu Leu Asp Leu Gly Arg Ser Ala Arg His His Val Ala
 325 330 335
 tcc ggc ttc ggg atc cac cag tgc ctg ggg cag aac ctc gcc cgc gcg 1475
 Ser Gly Phe Gly Ile His Gln Cys Leu Gly Gln Asn Leu Ala Arg Ala
 340 345 350
 gag atg gag atc gcc ctg cgc tca ctg ttc acc agg atc ccg cag ctg 1523
 Glu Met Glu Ile Ala Leu Arg Ser Leu Phe Thr Arg Ile Pro Gln Leu
 355 360 365
 cgc ctc gcc gtg cgc gcc gcc gag att cgc ttc aag gac gga gac acc 1571
 Arg Leu Ala Val Pro Ala Ala Glu Ile Pro Phe Lys Asp Gly Asp Thr
 370 375 380
 ctg caa ggc atg atc gaa ctg cgc ctg gcc tgg tag cagccaggac ggcaga 1623
 Leu Gln Gly Met Ile Glu Leu Pro Leu Ala Trp
 385 390 395
 ccaaagaaag gggccgga atg cgc atc ggc atc gac acc gac cgc tgt atc 1675
 Met Arg Ile Ala Ile Asp Thr Asp Arg Cys Ile
 400 405
 ggc gcc ggc cag tgt gcc ctg acc ggc ccc ggg ggt ttc acc cag gat 1723
 Gly Ala Gly Gln Cys Ala Leu Thr Ala Pro Gly Gly Phe Thr Gln Asp
 410 415 420
 gac gac ggt ttc agt gca ctg ctg ccc ggc cgc gag gac ggc gcc ggc 1771
 Asp Asp Gly Phe Ser Ala Leu Leu Pro Gly Arg Glu Asp Gly Ala Gly
 425 430 435
 gac cgc ctg gtg cgc gaa gcc gcc cgc gcc tgc ccc gtg cag gcc att 1819
 Asp Pro Leu Val Arg Glu Ala Ala Arg Ala Cys Pro Val Gln Ala Ile
 440 445 450
 ggc gtc acc gac gat tag cagcaccgcc gcggaccgac cggcagacgc gcgcggcc 1875
 Ala Val Thr Asp Asp
 455
 ccggctgaca ccggcgccc gaggcgcgcc cgagccgtcc gccctccac ttgtccctac 1935
 ggcattcacc ccatccgcta ccgcaacacc ccttgggiga cgggcagitt cgaggacccc 1995
 ggtctgcccg gggcgtactg gtgacctca ccggcttcac gccgcgaitg cccacatagg 2055
 cgtcgtcgtc cgggcgcac acgaagcgcg gtccgtgccc cggctcgtaa cggtcacaga 2115
 tggccggcag ttccacggg aaccgccggg ccacatcggg caccggggcc ggggccacca 2175
 acagggtgac cagcgtctc ctgcccgtc gcgcgacatc gttagagctg gcgaacagca 2235
 ccagcttgtc cggcgcaicc ggggaccgtc gcgccgccc ggcctgcggc gaggcaacct 2295
 tcagcgtcac cctcggcgcg cccaccacgt cgac 2329

<210> 3
 <211> 1860
 <212> DNA
 <213> Unknown

<220>
 <221> CDS
 <222> (172) .. (1383)
 <223> A-1560 strain

<220>
 <221> CDS
 <222> (1399) .. (1593)

<400> 3
 cggggaatcgt acgccgiacc gtttcggggc aaccgaattt cgaigcggaa tggatggttc 60
 ccagccagat ccgcaggta gccgatctgg ccgaacttga tctcgtcac tggatgcctc 120
 gggcatctaa tgaagatcgg cagcagcat ccttcgtcgc caggtctccc c atg aca 177
 Met Thr

EP 1 705 247 A1

5	gac acg aca gac ctg acc gag ctg tca gat ccc gtc tcc ttc ccc cag Asp Thr Thr 5 Asp Leu Thr Glu Leu Ser Asp Pro Val Ser Phe Pro Gln	225
	gac cgg agc tgc ccc tac cac ccc acc ggg tac gac ccc ctg cgc Asp Arg Ser Cys Pro Tyr His Pro Pro Thr Gly Tyr Asp Pro Leu Arg	273
10	acc gaa cgg ccc gcc cgc atc cgg ctg tac gac ggc cgc ccc gcc Thr Glu Arg Pro Pro Ala Arg Ile Arg Leu Tyr Asp Gly Arg Pro Ala	321
	tgg ctg gtc acc ggc cac gcc gtc gcc cgt gac ctg ctg gtc gac ccc Trp Leu Val Thr Gly His Ala Val Ala Arg Asp Leu Leu Val Asp Pro	369
15	cgc ctg tcc acg gac cgc acc cgc tgc ggc ttc ccc gcc aca act ccc Arg Leu Ser Thr Asp Arg Thr Arg Ser Gly Phe Pro Ala Thr Thr Pro	417
	cgc ttc gcc ggc gtc cgc gac cgc aag ccc ggc ctg ctg ggc gtc gac Arg Phe Ala Ala Val Arg Asp Arg Lys Pro Ala Leu Leu Gly Val Asp	465
20	gac ccc aag cac cgc acc cag cgg tgg atg atg atc ccc agc ttc acc Asp Pro Lys His Arg Thr Gln Arg Trp Met Met Ile Pro Ser Phe Thr	513
	ctc agg cgc gcc acc gag ctg agg ccc cgc atc cag gag atc gtc gac Leu Arg Arg Ala Thr Glu Leu Arg Pro Arg Ile Gln Glu Ile Val Asp	561
25	gaa ctg ctg gac gtg atg atc gcc cag gga ccc ccc gcc gac ctg gtg Glu Leu Leu Asp Val Met Ile Ala Gln Gly Pro Pro Ala Asp Leu Val	609
	cgt tcc ttc ggc ctg ccc gtg ccc tcc atg gtg atc tgc gcc ctg ctg Arg Ser Phe Ala Leu Pro Val Pro Ser Met Val Ile Cys Ala Leu Leu	657
30	ggc gtg ccc tac gcc gac cac gag ttc ttc gag gac cag tcc agg cgg Gly Val Pro Tyr Ala Asp His Glu Phe Phe Glu Asp Gln Ser Arg Arg	705
	ctg ctg cgc gga ccc gcc gcc gag gac acg cag gac gcc ccc gac cgc Leu Leu Arg Gly Pro Ala Ala Glu Asp Thr Gln Asp Ala Arg Asp Arg	753
35	ctc gcc cgc tac ctg gag gac ctg atc gac gag aag ccc cgc ccc ccc Leu Ala Ala Tyr Leu Glu Asp Leu Ile Asp Glu Lys Arg Arg Arg Pro	801
	ggt gac ggc ctg ctg gac gaa ctg gtc cag cag cgt ctg aac gaa ggc Gly Asp Gly Leu Leu Asp Glu Leu Val Gln Gln Arg Leu Asn Glu Gly	849
40	gag ctg gac cgc gag gaa ctg acc ggc ctg ggc atg atc ctg ctg gtc Glu Leu Asp Arg Glu Glu Leu Thr Ala Leu Ala Met Ile Leu Leu Val	897
	gcg ggc cac gag acc acc gcc aac atg atc tcc ctg ggc acc tac acg Ala Gly His Glu Thr Thr Ala Asn Met Ile Ser Leu Gly Thr Tyr Thr	945
45	ctc ctg ctg cac ccc gaa cgg ctg acc gag ctg cgc gcc gac ccc gcc Leu Leu Leu His Pro Glu Arg Leu Thr Glu Leu Arg Ala Asp Pro Ala	993
	ctg ctg ccc gcc gcc gtc gag gaa ctg atg cgg atg ctg tcc atc gcc Leu Leu Pro Ala Ala Val Glu Glu Leu Met Arg Met Leu Ser Ile Ala	1041
50	gac gga ctg ctg cgg cag gcc acc gag gac atc gag atc gcc ggg acc Asp Gly Leu Leu Arg Gln Ala Thr Glu Asp Ile Glu Ile Ala Gly Thr	1089
	acc atc agg gcc ggc gac ggc gtg gtc ttc tcc acc tct gtc atc aac Thr Ile Arg Ala Gly Asp Gly Val Val Phe Ser Thr Ser Val Ile Asn	1137
55	cgc gac gag gac gtc tac ccc gcc ccc gac acc ctg gac ttc cac cgc	1185

Arg Asp Glu Asp Val Tyr Pro Ala Pro Asp Thr Leu Asp Phe His Arg
 325 330 335
 5 tgc acc cgc cac cac gtc gcc ttc ggt ttc gga atc cac cag tgc ctc 1233
 Ser Thr Arg His His Val Ala Phe Gly Phe Gly Ile His Gln Cys Leu
 340 345 350
 ggc cag aac ctc gcc cgc acc gaa ctg gag atc gcc ctg cgc acg ctc 1281
 Gly Gln Asn Leu Ala Arg Thr Glu Leu Glu Ile Ala Leu Arg Thr Leu
 10 355 360 365 370
 ctc gaa cgg ctg ccc acg ctc cgg ctc gcc gcc cca cgg gag gaa atc 1329
 Leu Glu Arg Leu Pro Thr Leu Arg Leu Ala Ala Pro Pro Glu Glu Ile
 375 380 385
 ccc ttc aaa ccc gcc gac acc atc cag ggg atg ctg gaa ctc ccc gtc 1377
 Pro Phe Lys Pro Gly Asp Thr Ile Gln Gly Met Leu Glu Leu Pro Val
 15 390 395 400
 agc tgg taa gaggtgccg tc atg cat atc gag atc gac aag gac cgc tgc 1428
 Ser Trp Met His Ile Glu Ile Asp Lys Asp Arg Cys
 405 410
 atc ggc gcc gga cag tgc gcc ctg acc gcc cgg ggt gtg ttc acc cag 1476
 Ile Gly Ala Gly Gln Cys Ala Leu Thr Ala Pro Gly Val Phe Thr Gln
 20 420 425 430
 gac gac gac gcc ttc agt gac ctg ttg ccc gcc cgg gag gac gcc gcc 1524
 Asp Asp Asp Gly Phe Ser Asp Leu Leu Pro Gly Arg Glu Asp Gly Ala
 415 435 440 445
 ggc gac cgg atg gtc cgg gag gcc gcc agg gcc tgc ccc gtg agt gcc 1572
 Gly Asp Pro Met Val Arg Glu Ala Ala Arg Ala Cys Pro Val Ser Ala
 25 450 455 460
 atc acg ctg tcc gag gac ggg tag gggccgagc cgcgccgcc gccggtccgc 1626
 Ile Thr Leu Ser Glu Asp Gly
 465
 tgcgcggcgc ccgtgccgac gcggcggcgc gccggcccggt ccggtgccgc tgcgtgcc 1686
 ccgtggcccc ggcggcggt gattgactag ggtcccggg tgagcgaaca ggcccagaag 1746
 30 cctccgggg cccggccgc gaaagacacc gggacggcgc ccgggaaacc ccttcctcta 1806
 cgtcgtcgtc tgcgccgcgc gcatcgccga aggcgtcagc aagcigatca ccgc 1860

35 <210> 4
 <211> 29
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> STRANDNESS : single

40 <220>
 <223> TOPOLOGY : linear

45 <220>
 <223> Description of Artificial Sequence : 5Dm-3F Primer

45 <400> 4
 ttgcscctsc-csgtccctc satgtsat

29

50 <210> 5
 <211> 21
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> STRANDNESS : single

5 <220>
 <223> TOPOLOGY : linear
 <220>
 <223> Description of Artificial Sequence : 5Dm-3R Primer
 <400> 5
 10 gtigatsays gasgtsgaga a 21
 <210> 6
 <211> 30
 <212> DNA
 15 <213> Artificial Sequence
 <220>
 <223> STRANDNESS : single
 <220>
 20 <223> TOPOLOGY : linear
 <220>
 <223> Description of Artificial Sequence : 6PIN-2F Primer
 <400> 6
 25 gctgcgcctg gccctggagg acatcgagat 30
 <210> 7
 <211> 30
 <212> DNA
 30 <213> Artificial Sequence
 <220>
 <223> STRANDNESS : single
 <220>
 35 <223> TOPOLOGY : linear
 <220>
 <223> Description of Artificial Sequence : 6PIN-2R Primer
 <400> 7
 40 ctgttcctcg aagaactcgt ggtcggcgta 30
 <210> 8
 <211> 30
 45 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> STRANDNESS : single
 50 <220>
 <223> TOPOLOGY : linear
 <220>
 <223> Description of Artificial Sequence : DM-NdeF Primer
 55 <400> 8

gcccccatat gacggaactg acggacatca 30

5
 <210> 9
 <211> 30
 <212> DNA
 <213> Artificial Sequence

10
 <220>
 <223> STRANDNESS : single

<220>
 <223> TOPOLOGY : linear

15
 <220>
 <223> Description of Artificial Sequence : DM-SpeR Primer

<400> 9
 gggccactag tcagccggcc ggcticgtca 30

20
 <210> 10
 <211> 30
 <212> DNA
 <213> Artificial Sequence

25
 <220>
 <223> STRANDNESS : single

<220>
 <223> TOPOLOGY : linear

30
 <220>
 <223> Description of Artificial Sequence : DM-BglI Primer

<400> 10
 cgcatagatc ttcacccgag cgggigaica 30

35
 <210> 11
 <211> 30
 <212> DNA
 <213> Artificial Sequence

40
 <220>
 <223> STRANDNESS : single

<220>
 <223> TOPOLOGY : linear

45
 <220>
 <223> Description of Artificial Sequence : DM-BglR Primer

<400> 11
 tcccgagatc tgaaggicc gcgtcacgt 30

50
 <210> 12
 <211> 22
 <212> DNA
 <213> Artificial Sequence

55

```

5      <220>
      <223> STRANDNESS : single

      <220>
      <223> TOPOLOGY : linear

10     <220>
      <223> Description of Artificial Sequence : 5D-1R Primer

      <400> 12
      aggtgcccag cgagatcatg tt 22

15     <210> 13
      <211> 30
      <212> DNA
      <213> Artificial Sequence

20     <220>
      <223> STRANDNESS : single

      <220>
      <223> TOPOLOGY : linear

25     <220>
      <223> Description of Artificial Sequence : 7PIN-2F Primer

      <400> 13
      ccatgatcct gctggtggcc ggccatgaga 30

30     <210> 14
      <211> 30
      <212> DNA
      <213> Artificial Sequence

35     <220>
      <223> STRANDNESS : single

      <220>
      <223> TOPOLOGY : linear

40     <220>
      <223> Description of Artificial Sequence : 07-NdeF Primer

      <400> 14
      gccccatatg accgaagcca tcccctactt 30

45     <210> 15
      <211> 30
      <212> DNA
      <213> Artificial Sequence

50     <220>
      <223> STRANDNESS : single

      <220>
      <223> TOPOLOGY : linear

55     <220>
      <223> STRANDNESS : single

      <220>
      <223> TOPOLOGY : linear

```

<220>
 <223> Description of Artificial Sequence : 07-SpeR Primer

5 <400> 15
 gccactagtg ctaatcgicg gtgaccgcaa 30

10 <210> 16
 <211> 21
 <212> DNA
 <213> Artificial Sequence

15 <220>
 <223> STRANDNESS : single

<220>
 <223> TOPOLOGY : linear

20 <220>
 <223> Description of Artificial Sequence : 5Dm-2R Primer

<400> 16
 ctggatsgtg tcsccsgggt t 21

25 <210> 17
 <211> 30
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> STRANDNESS : single

<220>
 <223> TOPOLOGY : linear

35 <220>
 <223> Description of Artificial Sequence : 5PIN-2F Primer

<400> 17
 cggaaatccac cagtgcctcg gccagaacct 30

40 <210> 18
 <211> 30
 <212> DNA
 <213> Artificial Sequence

45 <220>
 <223> STRANDNESS : single

<220>
 <223> TOPOLOGY : linear

50 <220>
 <223> Description of Artificial Sequence : tpm-NdeF Primer

<400> 18
 ggccccatat gacagacacg acagacctga 30

<210> 19
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> STRANDNESS : single

<220>
 <223> TOPOLOGY : linear


<220>
 <223> Description of Artificial Sequence : tpm-SpeR Primer

<400> 19
 gcgcgactag tccccctacc cgtcctcgga


30

Claims

1. A DNA participating in biological transformation of a macrolide compound (hereinafter referred to as a macrolide compound 11107B) represented by the formula (I):



into a 16-position hydroxy macrolide compound represented by the formula (II):



the DNA being an isolated and pure DNA comprising a DNA encoding a protein having 16-position hydroxylating

enzymatic activity or ferredoxin, partly or entirely or its variant.

2. The DNA according to Claim 1, which is **characterized by** the following (a), (b) or (c):

(a) a DNA encoding a protein having the enzymatic activity in hydroxylating the 16-position of the macrolide compound 11107B and selected from the group consisting of a continuous nucleotide sequence from the base 1322 to base 2548 of the sequence No. 1; a continuous nucleotide sequence from the base 420 to base 1604 of the sequence No. 2; and a continuous nucleotide sequence from the base 172 to base 1383 of the sequence No. 3;

(b) a DNA which is a variant of the DNA described in the above (a);

(i) is hybridized with the DNA described in the above (a) under a stringent condition; and

(ii) encodes a protein having enzymatic activity in hydroxylating the 16-position of the macrolide compound 11107B; and

(c) a DNA encoding a protein having the same amino acid sequence as the protein encoded by the DNA described in the above (a) or (b) though it is not hybridized with the DNA described in the above (a) under a stringent condition because of the degeneracy of a gene codon.

3. A protein encoded by the DNA as claimed in Claim 2.

4. A self-replicative or integrating replicative recombinant plasmid carrying the DNA as claimed in Claim 2.

5. A transformant into which the recombinant plasmid as claimed in Claim 4 transforms.

6. A method of isolating a DNA encoding a protein having enzymatic activity in hydroxylating the 16-position of the macrolide compound 11107B, the method **characterized by** using the DNA as claimed in Claim 2 or a DNA constituted of a part of the DNA as a probe or a primer.

7. The DNA according to Claim 1, which is **characterized by** the following (d), (e) or (f):

(d) a DNA encoding ferredoxin and selected from the group consisting of a continuous nucleotide sequence from the base 2564 to base 2761 of the sequence No. 1, a continuous nucleotide sequence from the base 1643 to base 1834 of the sequence No. 2 and a continuous nucleotide sequence from the base 1399 to base 1593 of the sequence No. 3;

(e) a DNA which is a variant of the DNA represented by the above (d);

(i) is hybridized with the DNA described in the above (d) under a stringent condition; and

(ii) encodes a protein having a ferredoxin function; and

(f) a DNA encoding a protein having the same amino acid sequence as the protein encoded by the DNA represented by the above (d) or (e) though it is not hybridized with the DNA described in the above (d) under a stringent condition because of the degeneracy of a gene codon.


8. A protein encoded by the DNA as claimed in Claim 7.

9. A self-replicative or integrating replicative recombinant plasmid carrying the DNA as claimed in Claim 7.


10. A transformant into which the recombinant plasmid as claimed in Claim 9 transforms.

11. A method of isolating a DNA encoding a protein having a ferredoxin function, the method **characterized by** using the DNA as claimed in Claim 7 or a DNA constituted of a part of the DNA as a probe or a primer.

12. A method of producing a 16-position hydroxy macrolide compound, the method comprises the steps of culturing the transformant as claimed in Claim 5 or 10 in a medium; bringing the proliferated transformant into contact with a macrolide compound represented by the formula (III):



10 (wherein W represents



20 R^{12} , R^{16b} , R^{17a} , R^{17b} , R^{18} , R^{20a} , R^{20b} , R^{21a} and R^{21b} , which may be the same as or different from, respectively represent:

- 25 (1) hydrogen atom;
(2) a C_{1-22} alkyl group which may have a substituent;
(3) -OR (wherein R represents:

- 30 1) hydrogen atom; or
2) a C_{1-22} alkyl group;
3) a C_{7-22} aralkyl group;
4) a 5-membered to 14-membered heteroaryloxyalkyl group;
5) a C_{2-22} alkanoyl group;
6) a C_{7-15} aroyl group;
7) a C_{3-23} unsaturated alkanoyl group;
8) -COR^{oo} (wherein R^{oo} represents:

- 35 8-1) a 5-membered to 14-membered heteroaryloxyaryl group;
8-2) a C_{1-22} alkoxy group;
8-3) an unsaturated C_{2-22} alkoxy group;
8-4) a C_{6-14} aryloxy group;
40 8-5) a 5-membered to 14-membered heteroaryloxy group; or
8-6) a 3-membered to 14-membered nitrogen-containing non-aromatic heterocyclic group, each of which may have a substituent);

- 45 9) a C_{1-22} alkylsulfonyl group;
10) a C_{6-14} arylsulfonyl group; or
11) -SiR^{s1}R^{s2}R^{s3}, (wherein R^{s1}, R^{s2} and R^{s3}, which may be the same as or different from, respectively represent a C_{1-6} alkyl group or a C_{6-14} aryl group), each of which may have a substituent);

50 (4) a halogen atom; or

(5) -R^M-NR^{N1}R^{N2}, {wherein R^M represents a single bond or -O-CO-; and R^{N1} and R^{N2}

1) may be the same as or different from, respectively represent:

- 55 1-1) hydrogen atom; or
1-2)

- (i) a C_{1-22} alkyl group;
(ii) an unsaturated C_{2-22} alkyl group;


- (iii) a C₂₋₂₂ alkanoyl group;
- (iv) a C₇₋₁₅ aroyl group;
- (v) an unsaturated C₃₋₂₃ alkanoyl group;
- (vi) a C₆₋₁₄ aryl group;
- (vii) a 5-membered to 14-membered heteroaryl group;
- (viii) a C₇₋₂₂ aralkyl group;
- (ix) a C₁₋₂₂ alkylsulfonyl group; or
- (x) a C₆₋₁₄ arylsulfonyl group, each of which may have a substituent, or

2) and R^{N1} and R^{N2} may be combined with the nitrogen atom to which they bound, to form a 3-membered to 14-membered nitrogen-containing non-aromatic heterocyclic group], provided that R^{21a} and R^{21b} may be combined with each other to form (i) a ketone structure (=O) or (ii) an oxime structure (=NOR^{ox} (wherein R^{ox} represents a C₁₋₂₂ alkyl group, an unsaturated C₂₋₂₂ alkyl group, a C₆₋₁₄ aryl group, a 5-membered to 14-membered heteroaryl group or a C₇₋₂₂ aralkyl group, each of which may have a substituent));

R^{16a} represents hydrogen atom;

R^{21c} represents:


- (1) hydrogen atom; or
- (2)



(wherein R^{22a}, R^{22b} and R^{22c}, which may be the same as or different from, respectively represent:


- 1) hydrogen atom;
- 2) a C₁₋₆ alkyl group;
- 3) -OR (wherein R has the same meaning as the above);
- 4) -R^M-N^{R^{N1}}N^{R^{N2}} (wherein R^M, R^{N1} and R^{N2} have the same meanings as the above); or
- 5) a halogen atom, or

any one of R^{21a} and R^{21b} may be combined with any one of R^{22a} and R^{22b} to form the partial structure;



); and
G^m represents:

- (1) a group represented by the formula (GM-I):



{wherein

R² and R¹⁰, which may be the same as or different from, respectively represent hydrogen atom or a C₁₋₂₂ alkyl group;

R^{3a}, R^{3b}, R^{5a}, R^{5b}, R^{6a} and R^{6b}, which may be the same as or different from, respectively represent:

1) hydrogen atom;

2) hydroxyl group;

3)

3-1) a C₁₋₂₂ alkyl group;

3-2) a C₁₋₂₂ alkoxy group;

3-3) a C₆₋₁₄ aryloxy group;

3-4) a 5-membered to 14-membered heteroaryloxy group;

3-5) a C₂₋₂₂ alkanoyloxy group;

3-6) a C₇₋₁₅ aroyloxy group;

3-7) a C₃₋₂₃ unsaturated alkanoyloxy group;

3-8) -OCOR^{co} (wherein R^{co} has the same meaning as the above);

3-9) a C₁₋₂₂ alkylsulfonyloxy group;

3-10) a C₆₋₁₄ arylsulfonyloxy group; or

3-11) -OSIR^{s1}R^{s2}R^{s3} (wherein R^{s1}, R^{s2} and R^{s3} have the same meanings as the above), each of which may have a substituent;

4) a halogen atom; or


5) -R^M-NR^{N1}R^{N2} (wherein R^M, R^{N1} and R^{N2} have the same meanings as the above); or

R^{5a} and R^{5b} may be combined with each other to form a ketone structure (=O); or


R^{6a} and R^{6b} may be combined with each other to form a spirooxysilanyl group or an exomethylene group; or

R^{7a} and R^{7b}, which may be the same as or different from, respectively represent hydrogen atom or -OR^H (wherein R^H represents hydrogen atom, a C₁₋₂₂ alkyl group or a C₂₋₂₂ alkanoyl group));


(2) a group represented by the formula (GM-II) :




(wherein R², R^{3a}, R^{3b}, R^{6a}, R^{6b}, R^{7a}, R^{7b} and R¹⁰ have the same meanings as those in the formula (GM-I));
 (3) a group represented by the formula (GM-III):




(wherein R², R^{5a}, R^{5b}, R^{6a}, R^{6b}, R^{7a}, R^{7b} and R¹⁰ have the same meanings as those in the formula (GM-I));
 (4) a group represented by the formula (GM-IV):



(wherein R², R^{6a}, R^{7a}, R^{7b} and R¹⁰ have the same meanings as those in the formula (GM-I)); or
 (5) a group represented by the formula (GM-V):




(wherein R², R^{3a}, R^{6a}, R^{6b} and R¹⁰ have the same meanings as those in the formula (GM-I))) during or after
 culturing, to convert it into a 16-position hydroxy macrolide compound represented by the formula (IV):



(wherein W, R¹², R^{16b}, R^{17a}, R^{17b}, R^{20a}, R^{20b}, R^{21a}, R^{21b}, R^{21c} and G^m have the same meanings as those in the formula (III)); and then collecting the 16-position hydroxy macrolide compound thus converted.


13. A production method according to Claim 12, wherein the transformant is the transformant as claimed in Claim 5 and has a DNA encoding ferredoxin.
14. The production method according to Claim 12, the method comprises the step of converting a compound represented by the formula (III-a):




(wherein



represents a double bond or a single bond; W' represents a double bond or



; R^{5'} represents hydrogen atom or an acetoxy group; R^{6'} represents hydrogen atom or hydroxyl group; and R^{7'} represents hydrogen atom or acetyl group) into a compound represented by the formula (IV-a):



(wherein




5 , W' , R^{5'} , R^{6'} and R^{7'} have the same meanings as those in the formula (III-a)).

15. The production method according to Claim 14, wherein, in the conversion of the compound of the formula (III-a) into the compound of the formula (IV-a), the compound to be subjected is a compound selected from the group consisting of:

10

(1) a compound in which

15



20

is a single bond; W' is



25


and R^{5'} , R^{6'} and R^{7'} are respectively hydrogen atom;
(2) a compound in which

30



35

is a single bond, W' is



40

45


R^{5'} and R^{6'} are respectively hydrogen atom; and R^{7'} is acetyl group;
(3) a compound in which

50




55


is a single bond, W' is



R⁵ and R⁷ are respectively hydrogen atom; and R⁶ is hydroxyl group;
(4) a compound in which



is a single bond, W' is




R⁵ is hydrogen atom, R⁶ is hydroxy group; and R⁷ is acetyl group;
(5) a compound in which




is a single bond; W' is a double bond; and R⁵, R⁶ and R⁷ are respectively hydrogen atom;
(6) a compound in which




is a single bond; W' is a double bond; R⁵ and R⁶ are respectively hydrogen atom; and R⁷ is acetyl group;
(7) a compound in which



is a single bond; W' is a double bond; R⁵ and R⁷ are respectively hydrogen atom; and R⁶ is hydroxyl group;
(8) a compound in which




is a single bond; W' is a double bond; R⁵ is hydrogen atom; R⁶ is hydroxy group; and R⁷ is acetyl group;
(9) a compound in which



5

is a double bond; W' is


10



15

; R^{5'} and R^{7'} are respectively hydrogen atom; and R^{6'} is hydroxyl group;
(10) a compound in which

20



is a double bond; W' is


25



30

R^{5'} is hydrogen atom; R^{6'} is hydroxy group; and R^{7'} is acetyl group;
(11) a compound in which

35



40

is a single bond; W' is


45




50

R^{5'} is acetoxy group; R^{6'} is hydroxyl group; and R^{7'} is hydrogen atom; and
(12) a compound in which

55



is a single bond; W' is



5

R^{5'} is an acetoxy group; R^{6'} is hydroxyl group; and R^{7'} is acetyl group.

10 **16.** Use of the transformant as claimed in Claim 5 or 10 for producing a 16-position hydroxy macrolide compound.

15

20

25

30

35

40

45

50

55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2004/017906

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁷ C12N15/09, 1/15, 1/19, 1/21, 5/10, 9/02, C12P17/08, C07K14/00, C07D407/16 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl ⁷ C12N1/00-15/19, C12P17/08, C07K14/00-16/46, C07D407/16 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) SwissProt/PIR/GeneSeq, MEDLINE (STN), Genbank/EMBL/DBJ/GeneSeq, WPI (STN), BIOSIS (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X/ A	WO 2003/040370 A1 (SUMITOMO CHEMICAL CO., LTD.), 15 May, 2003 (15.05.03), Full text & JP 2004-57194 A & EP 1457558 A1	1-11/ 12-16
P,A	WO 2003/099813 A1 (MERCIAN Co.), 04 December, 2003 (04.12.03), (Family: none)	1-16
A	WO 2002/060890 A1 (MERCIAN CO.), 08 August, 2002 (08.08.02), (Family: none)	1-16
P,A	WO 2004/050890 A1 (MERCIAN CO.), 17 June, 2004 (17.06.04), (Family: none)	1-16
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 21 February, 2005 (21.02.05)		Date of mailing of the international search report 08 March, 2005 (08.03.05)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (January 2004)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2004/017906

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 2004/011459 A1 (MERCIAN CO.), 05 February, 2004 (05.02.04), (Family: none)	1-16
P, A	WO 2004/011661 A1 (MERCIAN CO.), 05 February, 2004 (05.02.04), (Family: none)	1-16
A	WO 2003/087381 A1 (MERCIAN CO.), 23 October, 2003 (23.10.03), & EP 1500704 A1	1-16

Form PCT/ISA/210 (continuation of second sheet) (January 2004)

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 02060890 A [0002]
- WO 03087381 A [0085] [0120] [0141]

Non-patent literature cited in the description

- Molecular Cloning. Current Protocols in Molecular Biology. John Wiley & Sons, 1987 [0032]
- *Cell Technology*, 1995, vol. 14, 591-593 [0075] [0110] [0131]